PMA
PERFORMANCE MONITORING FOR ACTION

JOHNS HOPKINS
BLOOMBERG SCHOOL
of PUBLIC HEALTH

BILL & MELINDA GATES INSTITUTE *for*
POPULATION *and* REPRODUCTIVE HEALTH

Democratic Republic of Congo (Kongo Central)
Snapshot of Indicators
Phase 2 (2021) Cross-sectional Survey

PMA www.pmadata.org   @PMA4Action   @pma4action   @PMA4Action

Background Characteristics of Respondents

Distribution of women age 15 to 49 by background characteristics.

Background Characteristics	Percentage of Sample	Weighted Number	Unweighted Number
Total			
All	100.0	1929	1929
Age Group			
15-19	20.0	385	424
20-24	18.0	346	347
25-29	15.9	307	296
30-34	13.9	267	254
35-39	13.2	254	251
40-44	10.4	201	194
45-49	8.7	168	163
Marital Status			
Married, or in union	60.6	1168	1117
Unmarried, sexually active	12.8	247	277
Parity			
0-1 children	15.9	306	321
2-3 children	26.4	510	486
4+ children	33.6	649	612
Education			
Never	9.2	178	164
Primary	27.4	528	479
Secondary	60.0	1158	1199
Higher	3.4	65	86
Wealth Quintile			
Lowest	17.3	334	235
Lower	17.4	336	286
Middle	19.4	375	322
Higher	22.3	430	479
Highest	23.5	454	607

Current Use of Contraception

Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics.

Background Characteristics	Any Method		Modern Method [1]		Traditional Method	
	All Women	Married [2] Women	All Women	Married [2] Women	All Women	Married [2] Women
Total						
All	40.9	46.9	26.3	27.8	14.7	19.1
Age Group						
15-19	21.8	31.0	14.6	16.9	7.3	14.0
20-24	51.1	50.0	35.9	30.4	15.2	19.5
25-29	51.8	53.9	32.9	32.5	18.8	21.4
30-34	50.6	53.8	32.7	34.0	17.9	19.8
35-39	46.2	50.5	27.5	28.9	18.8	21.6
40-44	41.2	44.9	24.3	24.6	16.8	20.3
45-49	20.2	22.4	11.3	12.3	8.9	10.1
Marital Status						
Married, or in union	46.9	na	27.8	na	19.1	na
Unmarried, sexually active	59.8	na	42.8	na	17.0	na
Parity						
0-1 children	42.8	40.6	30.8	23.3	12.1	17.2
2-3 children	49.1	52.3	33.6	34.6	15.5	17.7
4+ children	45.0	48.1	25.6	26.4	19.4	21.7
Education						
Never	36.4	42.4	18.0	20.4	18.4	22.0
Primary	35.2	41.3	19.8	23.1	15.4	18.2
Secondary	42.9	52.3	29.4	32.9	13.6	19.3
Higher	63.7	*34.6*	46.6	*24.6*	17.1	*10.0*
Wealth Quintile						
Lowest	41.3	50.2	29.7	35.4	11.6	14.9
Lower	36.6	42.8	19.8	21.8	16.8	21.0
Middle	35.9	41.7	21.7	23.3	14.2	18.4
Higher	41.2	48.3	24.6	23.5	16.6	24.8
Highest	47.7	52.7	33.9	36.1	13.8	16.6

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

2 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Contraception Method Mix

Percent distribution of current contraceptive users age 15 to 49, by method type, marital status, and background characteristics.

Background Characteristics	**All Users (N = 821)**										**Married Users[1] (N = 547)**									
	Sterilization	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtnl	Steriliztn	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtnl
Total																				
All	2.3	26.7	0.4	6.5	2.1	5.5	5.9	14.1	0.9	35.8	3.3	28.5	0.4	6.8	2.4	4.6	1.6	11.3	0.4	40.7
Age Group																				
15-19	0.0	11.1	0.0	7.5	0.0	9.4	13.2	22.8	2.8	33.3	--	--	--	--	--	--	--	--	--	--
20-24	0.0	25.4	0.0	2.3	2.0	6.2	8.1	26.1	0.2	29.7	0.0	26.4	0.0	3.3	3.4	2.8	0.9	24.0	0.0	39.0
25-29	0.6	31.8	0.0	3.8	2.8	3.8	5.9	14.3	0.5	36.4	0.8	32.0	0.0	4.2	1.8	5.0	2.8	13.4	0.3	39.7
30-34	3.9	29.1	0.0	12.1	3.4	4.0	2.9	8.9	0.3	35.4	4.6	31.3	0.0	10.0	3.6	3.1	0.9	9.5	0.0	36.9
35-39	0.6	29.7	1.5	6.9	1.6	7.9	3.1	6.9	1.2	40.6	0.8	28.5	0.9	7.8	1.9	8.5	1.9	6.9	0.2	42.7
40-44	9.0	27.4	1.4	9.0	2.4	3.8	4.8	0.0	1.4	40.9	10.0	23.2	1.5	9.9	2.7	3.9	2.1	0.0	1.5	45.2
45-49	*10.0*	*25.4*	*0.0*	*9.7*	*0.0*	*0.8*	*0.0*	*8.8*	*1.1*	*44.1*	*12.2*	*26.9*	*0.0*	*3.7*	*0.0*	*0.0*	*0.0*	*10.7*	*1.4*	*45.1*
Marital Status																				
Married, or in union	3.3	28.5	0.4	6.8	2.4	4.6	1.6	11.3	0.4	40.7	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	0.0	20.8	0.6	8.1	1.8	7.5	15.0	16.7	1.2	28.5	na	na	na	na	na	na	na	na	na	na
Parity																				
0-1 children	0.0	22.6	0.0	4.9	2.8	5.7	8.8	26.7	0.3	28.1	0.0	16.5	0.0	6.2	3.6	1.4	3.9	25.9	0.0	42.5
2-3 children	3.0	33.9	0.0	6.8	2.3	6.1	4.1	11.3	0.8	31.6	3.7	33.0	0.0	6.7	2.9	5.2	1.4	13.1	0.2	33.8
4+ children	3.6	30.9	1.0	7.5	1.8	4.5	0.5	6.5	0.6	43.2	3.8	29.1	0.7	7.1	1.9	4.4	0.6	6.5	0.6	45.2
Education																				
Never	3.2	25.4	0.0	1.6	2.8	10.1	0.0	6.2	0.2	50.5	3.6	24.9	0.0	0.0	3.2	9.6	0.0	6.9	0.0	51.8
Primary	3.9	29.8	0.5	5.3	1.5	5.5	0.6	9.2	0.1	43.7	4.2	31.3	0.5	4.7	1.7	3.7	0.7	9.0	0.1	44.2
Secondary	1.7	27.1	0.4	8.0	2.4	5.3	6.7	15.8	1.0	31.6	2.8	27.6	0.4	9.2	2.8	4.3	2.2	13.1	0.6	37.0
Higher	0.0	9.7	0.0	3.0	0.0	0.5	29.3	27.3	3.4	26.8	--	--	--	--	--	--	--	--	--	--
Wealth Quintile																				
Lowest	1.8	35.4	0.0	3.0	2.0	4.0	0.0	25.7	0.0	28.0	2.1	35.1	0.0	3.4	2.3	3.8	0.0	23.7	0.0	29.6
Lower	1.8	28.1	0.5	5.9	1.5	4.3	0.0	11.4	0.5	45.9	2.2	28.8	0.6	5.8	1.8	3.3	0.0	8.3	0.3	49.0
Middle	5.8	27.7	0.2	6.6	0.0	9.1	0.1	9.9	1.0	39.5	7.5	24.9	0.3	6.7	0.0	9.1	0.1	6.2	1.1	44.1
Higher	3.0	25.7	0.0	7.3	2.1	6.2	3.9	10.4	1.0	40.3	4.8	25.5	0.0	5.5	0.9	2.5	0.8	8.3	0.4	51.3
Highest	0.0	20.3	0.9	8.5	3.8	4.2	18.1	13.8	1.4	29.0	0.0	27.6	1.0	12.7	7.0	4.6	7.0	8.2	0.2	31.5

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Modern contraceptive methods

Percentage of women ages 15 to 49 using short-acting and long-acting contraceptive methods by background characteristics.

Background Characteristics	Short Acting Methods [1]		Long Acting Methods [2]	
	All Women (N = 1929)	Married Women [3] (N = 1117)	All Women (N = 1929)	Married Women [3] (N = 1117)
Total				
All	14.3	12.7	12.0	15.1
Age Group				
15-19	12.1	10.0	2.4	6.9
20-24	23.0	17.2	13.0	13.2
25-29	16.1	14.8	16.8	17.7
30-34	16.0	14.6	16.7	19.3
35-39	12.7	13.7	14.8	15.2
40-44	8.8	9.0	15.6	15.6
45-49	4.1	3.5	7.2	8.7
Marital Status				
Married, or in union	12.7	na	15.1	na
Unmarried, sexually active	30.0	na	12.8	na
Parity				
0-1 children	21.1	16.6	9.7	6.7
2-3 children	15.5	15.4	18.1	19.2
4+ children	9.6	10.2	16.0	16.2
Education				
Never	7.6	8.4	10.4	12.1
Primary	7.8	8.2	12.0	14.9
Secondary	16.8	16.9	12.5	16.0
Higher	40.4	*11.6*	6.2	*13.1*
Wealth Quintile				
Lowest	14.3	16.7	15.4	18.7
Lower	8.6	8.3	11.2	13.5
Middle	9.6	9.7	12.1	13.6
Higher	12.8	8.9	11.8	14.6
Highest	23.7	21.0	10.1	15.1

1   Short-acting methods include injectables, pills, emergency contraception, condom (male and female), standard days/cycle beads, and LAM.

2   Long-acting methods include IUD, implants, and sterilization (male and female).

3   Married includes married women and those who are living with a man

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Unmet Need for Family Planning

Percent of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics.

Background Characteristics	All Women (N = 1929)			Married Women[1] (N = 1117)		
	For Spacing	For Limiting	Total	For Spacing	For Limiting	Total
Total						
All	14.4	4.1	18.4	17.3	4.9	22.2
Age Group						
15-19	12.7	3.8	16.4	29.7	10.7	40.4
20-24	19.3	4.6	23.9	26.1	4.5	30.6
25-29	18.3	1.4	19.6	19.4	1.6	21.0
30-34	18.6	1.6	20.2	21.8	1.8	23.6
35-39	16.1	7.3	23.4	17.4	8.5	25.8
40-44	5.2	5.7	11.0	6.3	6.0	12.3
45-49	2.3	5.9	8.1	1.8	6.9	8.7
Marital Status						
Married, or in union	17.3	4.9	22.2	na	na	na
Unmarried, sexually active	22.6	6.0	28.6	na	na	na
Parity						
0-1 children	21.0	2.9	23.9	24.4	2.5	27.0
2-3 children	17.0	2.8	19.8	18.6	2.5	21.1
4+ children	13.2	6.8	20.0	13.9	7.0	20.9
Education						
Never	15.1	2.3	17.4	16.0	3.0	19.0
Primary	16.3	6.7	23.0	18.9	6.5	25.4
Secondary	13.8	3.4	17.2	16.6	4.4	21.0
Higher	6.8	0.0	6.8	*16.0*	*0.0*	*16.0*
Wealth Quintile						
Lowest	19.0	7.0	26.0	20.4	8.1	28.5
Lower	16.9	7.1	23.9	18.4	7.6	26.0
Middle	17.6	2.3	20.0	20.0	2.5	22.5
Higher	10.7	3.0	13.7	13.6	2.6	16.2
Highest	9.9	2.2	12.1	13.3	3.6	16.9

1   Married includes married women and those who are living with a man

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Need and Demand for Family Planning

Percent of women age 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics.

Background Characteristics	All Women (N=1929)					Married Women [2] (N=1117)				
	Any Method	Modern[1] Method	Besoins Unmet Need	Total Demand	Demand Satisfied By Modern	Any Method	Modern[1] Method	Besoins Unmet Need	Total Demand	Demand Satisfied By Modern
Total										
All	40.9	26.3	18.4	59.4	44.3	46.9	27.8	22.2	69.2	40.2
Age Group										
15-19	21.8	14.6	16.4	38.3	38.1	31.0	16.9	40.4	71.3	*23.7*
20-24	51.1	35.9	23.9	75.0	47.9	50.0	30.4	30.6	80.6	37.8
25-29	51.8	32.9	19.6	71.4	46.1	53.9	32.5	21.0	74.9	43.4
30-34	50.6	32.7	20.2	70.8	46.2	53.8	34.0	23.6	77.4	43.9
35-39	46.2	27.5	23.4	69.6	39.5	50.5	28.9	25.8	76.3	37.9
40-44	41.2	24.3	11.0	52.1	46.7	44.9	24.6	12.3	57.3	43.0
45-49	20.2	11.3	8.1	28.4	39.8	22.4	12.3	8.7	31.1	*39.5*
Marital Status										
Married, or in union	46.9	27.8	22.2	69.2	40.2	na	na	na	na	na
Unmarried, sexually active	59.8	42.8	28.6	88.4	48.4	na	na	na	na	na
Parity										
0-1 children	42.8	30.8	23.9	66.8	46.1	40.6	23.3	27.0	67.5	34.6
2-3 children	49.1	33.6	19.8	68.9	48.7	52.3	34.6	21.1	73.4	47.2
4+ children	45.0	25.6	20.0	65.0	39.3	48.1	26.4	20.9	69.0	38.2
Education										
Never	36.4	18.0	17.4	53.7	33.5	42.4	20.4	19.0	61.4	33.3
Primary	35.2	19.8	23.0	58.2	34.1	41.3	23.1	25.4	66.7	34.6
Secondary	42.9	29.4	17.2	60.1	48.8	52.3	32.9	21.0	73.2	44.9
Higher	63.7	46.6	6.8	70.5	66.1	*34.6*	*24.6*	*16.0*	*50.6*	*--*
Wealth Quintile										
Lowest	41.3	29.7	26.0	67.3	44.2	50.2	35.4	28.5	78.7	44.9
Lower	36.6	19.8	23.9	60.5	32.7	42.8	21.8	26.0	68.8	31.8
Middle	35.9	21.7	20.0	55.9	38.9	41.7	23.3	22.5	64.2	36.3
Higher	41.2	24.6	13.7	54.9	44.8	48.3	23.5	16.2	64.5	36.5
Highest	47.7	33.9	12.1	59.8	56.6	52.7	36.1	16.9	69.6	51.9

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method
2 Married includes married women and those who are living with a man

na Not applicable
-- Less than 25 unweighted observations
<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Pregnancy Intentions

Unintended births among women age 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics.

Background Characteristics	All Women with 1 or More Births (N=930)			Married[1] Women with 1 or More Births (N=732)		
	Wanted Then	Wanted Later	Not At All	Wanted Then	Wanted Later	Not At All
Total						
All	37.9	53.2	8.9	40.5	51.3	8.2
Age Group						
15-19	25.6	67.4	7.0	*39.0*	*46.4*	*14.5*
20-24	28.6	62.6	8.8	32.0	61.0	7.0
25-29	39.4	55.9	4.8	41.2	53.4	5.4
30-34	40.3	55.6	4.1	38.9	57.5	3.5
35-39	44.5	45.6	9.8	45.2	45.0	9.8
40-44	48.2	28.9	22.9	52.8	30.6	16.6
45-49	*37.2*	*30.0*	*32.8*	--	--	--
Marital Status						
Married, or in union	40.5	51.3	8.2	na	na	na
Unmarried, sexually active	24.7	68.1	7.2	na	na	na
Parity						
0-1 children	34.9	59.7	5.3	43.4	51.6	4.9
2-3 children	41.5	53.4	5.1	43.1	52.2	4.7
4+ children	36.3	49.7	14.0	37.6	50.4	12.0
Education						
Never	38.7	56.0	5.3	39.4	54.6	6.0
Primary	43.6	45.5	11.0	44.3	44.9	10.8
Secondary	34.1	57.4	8.6	37.4	55.6	7.0
Higher	--	--	--	--	--	--
Wealth Quintile						
Lowest	17.1	68.0	14.9	19.2	66.3	14.5
Lower	45.3	46.2	8.5	47.2	46.0	6.8
Middle	43.8	50.3	5.9	43.4	49.8	6.7
Higher	42.5	47.1	10.5	48.8	44.3	6.9
Highest	42.7	53.7	3.6	50.7	45.3	4.0

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Contraceptive Choice

Percent distribution of women age 15 to 49 who are currently using modern contraception[1] reporting on who decided on the contraceptive method, by background characteristics.

Background Characteristics	Self	Jointly with Partner/Provider	Provider/Partner Alone	(N = 441)
Total				
All	52.4	40.5	7.1	
Age Group				
15-19	69.2	20.4	10.5	
20-24	53.3	34.5	12.2	
25-29	37.5	56.2	6.3	
30-34	55.6	41.6	2.7	
35-39	60.7	34.2	5.1	
40-44	*41.9*	*56.1*	*2.0*	
45-49	*--*	*--*	*--*	
Marital Status				
Married, or in union	38.9	53.0	8.2	
Unmarried, sexually active	80.9	14.6	4.5	
Parity				
0-1 children	61.6	32.1	6.3	
2-3 children	46.9	44.0	9.1	
4+ children	47.0	48.0	5.0	
Education				
Never	*44.1*	*53.2*	*2.7*	
Primary	48.9	46.7	4.4	
Secondary	54.2	36.8	9.0	
Higher	*56.3*	*43.7*	*0.0*	
Wealth Quintile				
Lowest	*25.8*	*70.4*	*3.7*	
Lower	37.2	54.2	8.6	
Middle	49.9	42.8	7.3	
Higher	67.9	21.9	10.2	
Highest	66.2	27.9	5.8	

1   Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Method Information Index Components

Percentage of current modern contraceptive [1] users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics.

Background Characteristics	Informed about other methods [2]		Informed about side effects [2]		Told what to do if experienced side effects [4]		Told about switching methods in future [2]	
	All Users	Married [3] Users	All Users	Married [3] Users	All Users	Married [3] Users	All Users	Married [3] Users
Total	**(441)**	**(274)**	**(441)**	**(274)**	**(222)**	**(164)**	**(441)**	**(274)**
All	38.3	40.6	54.8	64.1	76.0	75.1	39.9	40.1
Age Group								
15-19	27.7	--	25.4	--	--	--	25.4	--
20-24	40.2	55.5	50.7	62.0	55.3	48.5	42.7	46.5
25-29	31.2	32.2	51.1	57.2	71.3	67.2	29.8	29.5
30-34	46.9	40.7	57.1	61.9	81.8	81.2	46.3	40.8
35-39	39.0	35.7	71.4	75.0	88.1	86.5	45.6	41.1
40-44	40.4	39.8	70.4	71.4	82.8	83.2	48.6	52.3
45-49	--	--	--	--	--	--	--	--
Marital Status								
Married, or in union	40.6	na	64.1	na	75.1	na	40.1	na
Unmarried, sexually active	27.7	na	27.6	na	86.5	na	39.9	na
Parity								
0-1 children	46.7	72.2	47.1	55.1	71.3	--	43.0	52.9
2-3 children	40.9	42.1	51.6	57.0	73.8	74.1	38.3	34.4
4+ children	35.6	31.9	72.9	71.6	80.7	79.7	42.6	42.1
Education								
Never	26.6	--	71.7	--	--	--	35.2	--
Primary	38.1	35.2	55.9	57.9	70.1	68.4	36.4	33.0
Secondary	38.9	45.8	53.4	66.5	75.8	73.2	41.9	44.6
Higher	47.1	--	45.5	--	--	--	35.9	--
Wealth Quintile								
Lowest	39.9	32.8	83.2	80.0	69.5	68.5	26.2	26.3
Lower	40.5	41.7	56.7	61.7	62.6	61.4	37.8	37.0
Middle	30.9	32.1	50.0	50.0	88.7	--	32.9	34.1
Higher	36.2	46.5	49.9	63.3	79.8	76.2	47.4	51.1
Highest	41.9	49.9	41.6	59.1	81.7	82.2	47.6	52.9

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

2 Asked of all current/recent modern contraceptive users

3 Married includes married women and those who are living with a man

4 Asked of all current/recent modern contraceptive users who were informed about side effects

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Quality of Care

Percentage of current modern[1] contraceptive users who would return, would refer a friend/relative, or would return and refer a friend/relative to provider, by background characteristics.

Background Characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Total	**(N = 433)**	**(N = 433)**	**(N = 433)**
All	63.4	46.7	38.1
Age Group			
15-19	39.3	35.2	25.8
20-24	64.1	45.4	38.7
25-29	72.3	52.1	42.3
30-34	67.3	38.3	33.2
35-39	60.0	40.6	34.7
40-44	*67.5*	*74.7*	*54.0*
45-49	--	--	--
Marital Status			
Married, or in union	64.8	48.8	39.6
Unmarried, sexually active	59.8	44.5	35.2
Parity			
0-1 children	69.6	51.2	44.9
2-3 children	57.0	44.5	30.7
4+ children	69.0	49.7	43.1
Education			
Never	*71.3*	*48.5*	*39.2*
Primary	62.2	50.9	40.8
Secondary	63.1	44.7	37.1
Higher	*62.1*	*51.2*	*37.8*
Wealth Quintile			
Lowest	*82.1*	*65.7*	*65.7*
Lower	62.5	29.2	24.6
Middle	52.2	43.4	27.5
Higher	63.8	46.4	38.7
Highest	57.7	44.9	32.2

1   Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Knowledge of Family Planning

Percentage of women age 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics.

Background Characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All Women	Married[1] Women	All Women	Married[1] Women	All Women	Married[1] Women
Total	**(1929)**	**(1117)**	**(1929)**	**(1117)**	**(1929)**	**(1117)**
All	9.5	10.3	8.8	10.7	14.9	16.5
Age Group						
15-19	7.6	11.2	3.6	7.0	9.5	12.3
20-24	10.2	10.3	9.8	12.2	16.0	16.3
25-29	9.2	8.2	11.0	10.8	16.8	16.0
30-34	12.0	11.3	12.6	13.2	20.0	19.7
35-39	11.5	12.9	14.6	17.2	21.2	23.8
40-44	8.0	9.5	5.9	5.7	11.0	11.7
45-49	8.0	8.5	3.2	2.3	9.0	9.2
Marital Status						
Married, or in union	10.3	na	10.7	na	16.5	na
Unmarried, sexually active	9.9	na	7.4	na	14.6	na
Parity						
0-1 children	9.1	10.6	8.3	10.3	13.7	15.6
2-3 children	10.4	10.6	11.8	11.8	18.3	18.3
4+ children	9.9	10.6	10.5	10.7	16.0	16.6
Education						
Never	5.2	6.1	6.8	7.9	9.8	11.2
Primary	9.4	10.5	9.1	10.7	14.0	15.6
Secondary	9.7	10.9	9.3	11.6	15.7	18.2
Higher	19.8	*16.1*	3.2	*5.8*	22.9	*21.9*
Wealth Quintile						
Lowest	11.4	13.1	8.8	10.5	16.7	18.8
Lower	7.7	8.9	9.5	11.7	13.6	15.6
Middle	7.1	6.7	10.2	11.0	13.6	14.4
Higher	8.7	10.2	7.7	8.9	12.1	13.4
Highest	12.2	12.8	8.3	11.4	18.6	20.9

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Response Rates

Response rates of households and individuals, by residence.

Result	Urban	Rural	Total
Household Interviews			
Households Selected	N/A	N/A	2026
Households Occupied	N/A	N/A	1918
Households Interviewed	N/A	N/A	1891
Household Response Rate [1]	0.00%	0.00%	98.59%
Interviews with Women			
Number of Eligible Women [2]	N/A	N/A	1962
Number of Eligible Women Interviewed	N/A	N/A	1929
Eligible Woman Response Rate [3]	0.00%	0.00%	98.32%

1 *Household response rate = households interviewed/households selected*

2 *Eligible women are women ages 15-49 identified in completed household interviews*

3 *Eligible response rate = eligible women interviewed/eligible women*

Sample Error Estimates

Variable	All Women				Married Women[1]			
	Mean	Std Error	CI Lower	CI Upper	Mean	Std Error	CI Lower	CI Upper
Currently using a modern method	0.263	0.025	0.217	0.315	0.278	0.032	0.219	0.347
Currently using a traditional method	0.147	0.017	0.116	0.183	0.191	0.022	0.152	0.238
Currently using any contraceptive method	0.409	0.030	0.352	0.469	0.469	0.039	0.392	0.548
Currently using DMPA-IM	0.065	0.012	0.044	0.095	0.068	0.014	0.044	0.102
Currently using DMPA-SC	0.021	0.008	0.010	0.043	0.024	0.010	0.011	0.054
Currently using condoms	0.141	0.029	0.092	0.209	0.113	0.031	0.065	0.189
Currenlty using implants	0.267	0.031	0.210	0.332	0.285	0.034	0.223	0.358
Chose method by self or jointly in past 12 months	0.929	0.017	0.888	0.956	0.918	0.022	0.863	0.953
Informed by provider about other methods	0.383	0.037	0.311	0.460	0.406	0.035	0.338	0.479
Informed by provider about side effects	0.548	0.052	0.443	0.649	0.641	0.044	0.548	0.724
Satisfied with provider: Would return and refer friend/relative to provider	0.381	0.057	0.274	0.500	0.396	0.062	0.280	0.525
Visited by health worker who talked about family planning in past 12 months	0.095	0.020	0.062	0.144	0.103	0.024	0.064	0.160

1   Married includes married women and those who are living with a man