PMA
PERFORMANCE MONITORING FOR ACTION

JOHNS HOPKINS BLOOMBERG SCHOOL of PUBLIC HEALTH | BILL & MELINDA GATES INSTITUTE *for* POPULATION *and* REPRODUCTIVE HEALTH

Democratic Republic of Congo (Kinshasa)
Snapshot of Indicator
Phase 1 (2020) Survey

PMA www.pmadata.org   @PMA4Action   @pma4action   @PMA4Action

Background Characteristics of Respondents

Distribution of women age 15 to 49 by background characteristics.

Background Characteristics	Percentage of Sample	Weighted Number	Unweighted Number
Total			
All	100.0	2611	2611
Age Group			
15-19	22.2	580	573
20-24	20.9	546	543
25-29	16.3	426	429
30-34	12.3	322	340
35-39	10.7	279	281
40-44	10.4	271	266
45-49	7.2	187	179
Marital Status			
Married, or in union	42.7	1114	1159
Unmarried, sexually active	20.3	530	482
Parity			
0-1 children	56.8	1484	1472
2-3 children	23.9	625	629
4+ children	19.2	502	510
Residence			
Urban	100.0	2611	2611
Education			
Never	0.4	10	10
Primary	7.4	193	200
Secondary	72.9	1904	1903
Higher	19.3	505	498
Wealth Quintile			
Lowest	18.7	489	513
Lower	18.9	493	551
Middle	20.1	524	498
High	20.2	528	530
Highest	22.1	577	519
Region			
Kinshasa	100.0	2611	2611

Current Use of Contraception

Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics.

Background Characteristics	Any Method		Modern Method [1]		Traditional Method	
	All Women	Married [2] Women	All Women	Married [2] Women	All Women	Married [2] Women
Total						
All	43.6	54.7	24.5	29.6	19.1	25.0
Age Group						
15-19	21.7	*40.2*	11.6	*17.5*	10.2	*22.7*
20-24	47.3	49.5	28.6	36.8	18.7	12.7
25-29	57.3	53.7	35.6	30.2	21.7	23.5
30-34	60.8	67.5	35.8	39.5	25.0	28.0
35-39	53.1	64.6	26.2	31.6	26.9	33.0
40-44	44.2	50.7	20.1	21.9	24.1	28.8
45-49	24.8	30.0	11.7	14.9	13.2	15.0
Marital Status						
Married, or in union	54.7	na	29.6	na	25.0	na
Unmarried, sexually active	72.7	na	42.3	na	30.4	na
Parity						
0-1 children	35.5	37.2	20.5	22.0	15.1	15.2
2-3 children	52.7	58.2	30.5	32.5	22.2	25.7
4+ children	56.1	61.7	28.9	31.3	27.3	30.4
Residence						
Urban	43.6	54.7	24.5	29.6	19.1	25.0
Education						
Never	--	--	--	--	--	--
Primary	47.4	60.2	31.6	41.9	15.9	18.3
Secondary	41.7	54.9	22.4	27.9	19.3	26.9
Higher	49.3	50.6	29.6	29.1	19.7	21.5
Wealth Quintile						
Lowest	46.9	57.2	28.1	34.1	18.8	23.1
Lower	41.4	55.6	23.2	32.3	18.2	23.4
Middle	43.5	52.7	20.5	23.4	23.0	29.3
High	42.6	50.8	23.9	24.8	18.7	25.9
Highest	43.8	57.3	26.6	33.9	17.2	23.3
Region						
Kinshasa	43.6	54.7	24.5	29.6	19.1	25.0

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactional amenorrhea method

2 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Contraception Method Mix

Percent distribution of current contraceptive users age 15 to 49, by method type, marital status, and background characteristics.

Background Characteristics	All Users (N = 1139)										Married Users[1] (N = 609)									
	Sterilization	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtnl	Steriliztn	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtnl
Total																				
All	1.1	15.3	0.3	4.6	1.8	2.7	13.2	15.6	1.6	43.9	1.9	21.7	0.6	5.8	2.8	3.2	6.8	10.0	1.3	45.8
Age Group																				
15-19	0.0	6.9	0.0	1.3	0.0	2.4	16.1	23.0	3.5	46.8	--	--	--	--	--	--	--	--	--	--
20-24	0.0	12.7	0.0	2.6	2.6	2.4	18.5	20.1	1.7	39.6	0.0	30.4	0.0	4.7	9.5	4.0	11.3	14.5	0.0	25.7
25-29	0.0	12.8	0.0	8.4	1.6	2.7	17.3	17.9	1.4	37.9	0.0	19.7	0.0	9.7	1.7	3.4	10.4	10.1	1.4	43.7
30-34	0.2	20.9	0.5	7.1	2.3	1.9	12.6	10.9	2.5	41.2	0.3	23.9	0.6	6.9	2.9	2.4	8.7	9.7	3.2	41.4
35-39	1.3	22.5	2.0	3.3	1.1	4.2	5.1	8.9	0.9	50.7	1.5	21.5	2.4	3.8	1.3	4.9	3.6	8.8	1.1	51.1
40-44	4.1	17.4	0.0	2.6	1.8	2.4	4.7	12.5	0.0	54.5	4.7	16.2	0.0	2.9	2.1	2.4	4.0	11.0	0.0	56.7
45-49	*10.5*	*14.7*	*0.0*	*3.0*	*3.3*	*4.4*	*3.5*	*7.5*	*0.0*	*53.1*	*13.0*	*19.1*	*0.0*	*3.9*	*4.3*	*2.2*	*0.0*	*7.3*	*0.0*	*50.2*
Marital Status																				
Married, or in union	1.9	21.7	0.6	5.8	2.8	3.2	6.8	10.0	1.3	45.8	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	0.0	6.9	0.0	3.7	0.6	2.6	18.8	24.2	1.5	41.8	na	na	na	na	na	na	na	na	na	na
Parity																				
0-1 children	0.0	5.3	0.0	3.1	0.4	2.3	20.0	24.1	2.3	42.4	0.0	15.2	0.0	7.6	0.0	3.6	14.5	16.7	1.6	40.9
2-3 children	1.8	23.0	0.3	6.5	2.3	1.7	11.9	8.8	1.5	42.2	2.2	22.0	0.4	6.1	2.7	1.8	8.9	9.8	2.0	44.1
4+ children	2.2	25.1	1.1	5.0	3.7	4.5	1.8	7.6	0.5	48.6	2.4	23.9	1.2	4.8	4.0	4.5	1.8	7.6	0.5	49.3
Residence																				
Urban	1.1	15.3	0.3	4.6	1.8	2.7	13.2	15.6	1.6	43.9	1.9	21.7	0.6	5.8	2.8	3.2	6.8	10.0	1.3	45.8
Education																				
Never	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Primary	0.3	37.4	0.0	11.2	4.1	6.7	4.7	1.6	0.5	33.5	0.4	39.6	0.0	11.7	4.0	7.2	5.2	0.8	0.7	30.4
Secondary	1.4	16.0	0.5	4.6	1.9	2.6	12.5	12.9	1.3	46.2	2.5	20.9	0.9	5.2	2.9	2.5	5.8	9.0	1.2	49.1
Higher	0.2	4.9	0.0	2.1	0.6	1.2	18.6	29.5	3.0	39.9	0.6	11.6	0.0	3.5	1.6	2.7	12.9	22.0	2.6	42.4
Wealth Quintile																				
Lowest	0.6	27.9	0.5	6.2	2.5	5.0	8.4	6.4	2.3	40.1	1.2	33.9	1.1	7.6	3.3	6.7	3.6	2.3	0.0	40.3
Lower	1.1	18.8	0.7	6.3	2.2	4.5	10.5	11.3	0.8	43.8	2.1	27.4	1.3	9.3	4.1	2.9	4.4	5.2	1.4	42.0
Middle	1.1	10.8	0.0	2.0	1.1	2.2	10.7	18.0	1.3	52.9	2.1	16.8	0.0	2.9	1.3	3.6	5.7	10.5	1.6	55.6
High	1.0	8.3	0.0	6.1	1.6	1.0	17.7	18.5	2.1	43.8	1.8	13.0	0.0	8.7	2.9	1.5	6.9	12.5	1.7	51.1
Highest	1.5	11.5	0.5	2.7	1.6	1.1	17.9	22.6	1.5	39.2	2.4	18.9	0.9	1.8	2.7	1.8	11.9	17.2	1.7	40.7
Region																				
Kinshasa	1.1	15.3	0.3	4.6	1.8	2.7	13.2	15.6	1.6	43.9	1.9	21.7	0.6	5.8	2.8	3.2	6.8	10.0	1.3	45.8

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Modern contraceptive methods

Percentage of women ages 15 to 49 using short-acting and long-acting contraceptive methods by background characteristics.

Background Characteristics	Short Acting Methods [1]		Long Acting Methods [2]	
	All Women (N = 2611)	Married Women[3] (N = 1114)	All Women (N = 2611)	Married Women[3] (N = 1114)
Total				
All	17.2	16.4	7.3	13.3
Age Group				
15-19	10.1	*3.8*	1.5	*13.7*
20-24	22.6	21.7	6.0	15.0
25-29	28.2	19.6	7.3	10.6
30-34	22.6	22.8	13.1	16.8
35-39	12.5	15.2	13.7	16.4
40-44	10.6	11.4	9.5	10.6
45-49	5.4	5.3	6.3	9.6
Marital Status				
Married, or in union	16.4	na	13.3	na
Unmarried, sexually active	37.3	na	5.0	na
Parity				
0-1 children	18.6	16.3	1.9	5.6
2-3 children	17.3	18.3	13.2	14.3
4+ children	13.0	14.4	15.9	16.9
Residence				
Urban	17.2	16.4	7.3	13.3
Education				
Never	--	--	--	--
Primary	13.7	17.8	17.9	24.1
Secondary	14.9	14.6	7.5	13.3
Higher	27.1	23.0	2.5	6.2
Wealth Quintile				
Lowest	14.5	13.4	13.6	20.7
Lower	14.7	15.1	8.6	17.1
Middle	15.3	13.5	5.1	9.9
High	20.0	17.3	4.0	7.5
Highest	20.7	21.3	5.9	12.7
Region				
Kinshasa	17.2	16.4	7.3	13.3

1 Short-acting methods include injectables, pills, emergency contraception, condom (male and female), standard days/cycle beads, and LAM.

2 Long-acting methods include IUD, implants, and sterilization (male and female).

3 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Unmet Need for Family Planning

Percent of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics.

Background Characteristics	All Women (N = 2611)			Married Women[1] (N = 1114)		
	For Spacing	For Limiting	Total	For Spacing	For Limiting	Total
Total						
All	8.5	2.2	10.7	11.4	4.2	15.6
Age Group						
15-19	6.6	0.4	7.1	*9.2*	*1.9*	*11.2*
20-24	10.9	0.4	11.3	18.9	1.2	20.1
25-29	14.0	0.9	14.9	21.4	1.6	23.1
30-34	8.7	2.2	10.9	9.4	2.7	12.1
35-39	8.0	3.3	11.2	11.4	3.0	14.4
40-44	4.7	6.8	11.5	4.9	8.3	13.1
45-49	0.6	7.3	7.9	0.4	10.1	10.5
Marital Status						
Married, or in union	11.4	4.2	15.6	na	na	na
Unmarried, sexually active	15.4	1.6	17.0	na	na	na
Parity						
0-1 children	7.3	0.2	7.5	12.5	0.2	12.7
2-3 children	10.9	2.3	13.1	11.7	2.8	14.5
4+ children	9.2	7.8	17.0	10.4	8.1	18.5
Residence						
Urban	8.5	2.2	10.7	11.4	4.2	15.6
Education						
Never	--	--	--	--	--	--
Primary	12.5	2.8	15.3	14.6	3.4	18.0
Secondary	9.2	2.4	11.6	11.8	4.6	16.4
Higher	4.5	0.7	5.2	7.8	1.7	9.5
Wealth Quintile						
Lowest	13.0	2.4	15.5	17.1	4.9	22.0
Lower	11.6	2.8	14.4	13.6	4.9	18.5
Middle	9.0	2.2	11.2	13.3	5.2	18.4
High	6.4	1.9	8.3	10.5	2.7	13.1
Highest	3.6	1.6	5.2	4.4	3.7	8.1
Region						
Kinshasa	8.5	2.2	10.7	11.4	4.2	15.6

1 *Married includes married women and those who are living with a man*

na *Not applicable*

-- *Less than 25 unweighted observations*

<i> *Italics indicate estimate was generated from 25 to 49 unweighted observations*

Need and Demand for Family Planning

Percent of women age 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics.

Background Characteristics	All Women (N=2611)					Married Women [2] (N=1114)				
	Any Method	Modern[1] Method	Unmet Need	Total Demand	Demand Satisfied By Modern	Any Method	Modern[1] Method	Unmet Need	Total Demand	Demand Satisfied By Modern
Total										
All	43.6	24.5	10.7	54.3	45.1	54.7	29.6	15.6	70.3	42.2
Age Group										
15-19	21.7	11.6	7.1	28.8	40.1	*40.2*	*17.5*	*11.2*	*51.4*	*--*
20-24	47.3	28.6	11.3	58.6	48.8	49.5	36.8	20.1	69.7	52.8
25-29	57.3	35.6	14.9	72.2	49.2	53.7	30.2	23.1	76.7	39.4
30-34	60.8	35.8	10.9	71.7	49.9	67.5	39.5	12.1	79.7	49.6
35-39	53.1	26.2	11.2	64.4	40.7	64.6	31.6	14.4	79.0	40.0
40-44	44.2	20.1	11.5	55.7	36.1	50.7	21.9	13.1	63.8	34.4
45-49	24.8	11.7	7.9	32.8	35.6	30.0	14.9	10.5	40.5	*36.9*
Marital Status										
Married, or in union	54.7	29.6	15.6	70.3	42.2	na	na	na	na	na
Unmarried, sexually active	72.7	42.3	17.0	89.7	47.1	na	na	na	na	na
Parity										
0-1 children	35.5	20.5	7.5	43.0	47.5	37.2	22.0	12.7	49.9	44.1
2-3 children	52.7	30.5	13.1	65.8	46.3	58.2	32.5	14.5	72.7	44.7
4+ children	56.1	28.9	17.0	73.1	39.5	61.7	31.3	18.5	80.2	39.0
Residence										
Urban	43.6	24.5	10.7	54.3	45.1	54.7	29.6	15.6	70.3	42.2
Education										
Never	--	--	--	--	--	--	--	--	--	--
Primary	47.4	31.6	15.3	62.7	50.3	60.2	41.9	18.0	78.2	53.6
Secondary	41.7	22.4	11.6	53.3	42.1	54.9	27.9	16.4	71.2	39.2
Higher	49.3	29.6	5.2	54.4	54.4	50.6	29.1	9.5	60.1	48.5
Wealth Quintile										
Lowest	46.9	28.1	15.5	62.4	45.1	57.2	34.1	22.0	79.2	43.1
Lower	41.4	23.2	14.4	55.8	41.7	55.6	32.3	18.5	74.1	43.6
Middle	43.5	20.5	11.2	54.6	37.5	52.7	23.4	18.4	71.1	32.9
High	42.6	23.9	8.3	50.9	47.0	50.8	24.8	13.1	63.9	38.9
Highest	43.8	26.6	5.2	49.0	54.3	57.3	33.9	8.1	65.4	51.9
Region										
Kinshasa	43.6	24.5	10.7	54.3	45.1	54.7	29.6	15.6	70.3	42.2

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactional amenorrhea method

2 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Pregnancy Intentions

Unintended births among women age 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics.

Background Characteristics	All Women with 1 or More Births (N=1010)			Married[1] Women with 1 or More Births (N=718)		
	Wanted Then	Wanted Later	Not At All	Wanted Then	Wanted Later	Not At All
Total						
All	46.7	48.3	5.0	52.1	42.1	5.8
Age Group						
15-19	21.0	71.5	7.4	--	--	--
20-24	31.6	65.8	2.6	50.7	47.3	2.0
25-29	47.4	51.0	1.7	49.7	48.3	2.0
30-34	55.5	41.8	2.7	55.3	41.3	3.4
35-39	56.9	37.2	5.9	55.4	37.9	6.6
40-44	59.4	25.6	15.0	53.0	29.5	17.5
45-49	--	--	--	--	--	--
Marital Status						
Married, or in union	52.1	42.1	5.8	na	na	na
Unmarried, sexually active	30.3	67.8	1.9	na	na	na
Parity						
0-1 children	42.8	54.2	2.9	65.3	33.7	1.0
2-3 children	54.1	44.6	1.3	59.0	39.3	1.7
4+ children	40.3	47.3	12.3	35.9	50.4	13.7
Residence						
Urban	46.7	48.3	5.0	52.1	42.1	5.8
Education						
Never	--	--	--	--	--	--
Primary	35.3	53.5	11.2	38.0	49.6	12.4
Secondary	44.8	51.0	4.2	51.2	44.1	4.8
Higher	65.3	31.3	3.4	67.3	28.9	3.9
Wealth Quintile						
Lowest	32.5	61.1	6.4	34.8	56.8	8.5
Lower	40.5	53.4	6.1	44.6	48.9	6.5
Middle	44.5	49.1	6.3	50.6	42.6	6.8
High	57.5	39.9	2.6	63.2	33.2	3.6
Highest	62.5	34.2	3.3	66.0	30.0	4.0
Region						
Kinshasa	46.7	48.3	5.0	52.1	42.1	5.8

1 *Married includes married women and those who are living with a man*

na *Not applicable*

-- *Less than 25 unweighted observations*

<i> *Italics indicate estimate was generated from 25 to 49 unweighted observations*

Contraceptive Choice

Percent distribution of women age 15 to 49 who are currently using modern contraception[1] reporting on who decided on the contraceptive method, by background characteristics.

Background Characteristics	Self	Jointly with Partner/Provider	Provider/Partner Alone	(N = 593)
Total				
All	48.3	38.7	13.1	
Age Group				
15-19	35.2	47.9	16.9	
20-24	58.6	26.8	14.7	
25-29	51.0	39.6	9.3	
30-34	54.9	36.3	8.8	
35-39	44.1	39.6	16.3	
40-44	*25.6*	*52.5*	*21.9*	
45-49	--	--	--	
Marital Status				
Married, or in union	49.2	40.6	10.2	
Unmarried, sexually active	46.2	36.6	17.2	
Parity				
0-1 children	45.5	38.1	16.4	
2-3 children	55.4	34.3	10.4	
4+ children	44.4	45.5	10.1	
Residence				
Urban	48.3	38.7	13.1	
Education				
Never	--	--	--	
Primary	64.5	31.9	3.6	
Secondary	49.1	36.7	14.2	
Higher	39.2	46.8	14.0	
Wealth Quintile				
Lowest	63.2	24.2	12.7	
Lower	53.9	34.4	11.7	
Middle	45.9	42.7	11.4	
High	45.9	38.6	15.5	
Highest	34.1	52.2	13.7	
Region				
Kinshasa	48.3	38.7	13.1	

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Method Information Index Components

Percentage of current modern contraceptive [1] users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics.

Background Characteristics	Informed about other [2] methods		Informed about side [2] effects		Told what to do if experienced [2] side effects		Told about switching [2] methods in future	
	All Users	Married [3] Users	All Users	Married [3] Users	All Users	Married [3] Users	All Users	Married [3] Users
Total	**(596)**	**(320)**	**(596)**	**(320)**	**(227)**	**(163)**	**(596)**	**(320)**
All	32.6	43.8	40.9	53.5	81.3	83.9	-0.1	44.8
Age Group								
15-19	17.6	--	24.8	--	--	--	-107.1	--
20-24	23.2	*31.0*	31.2	*47.1*	*90.8*	--	-67.1	*35.6*
25-29	30.3	40.4	37.8	53.1	69.6	*80.6*	33.4	42.9
30-34	37.5	42.5	53.1	55.0	78.0	*80.8*	43.4	45.0
35-39	40.3	39.0	52.7	52.6	*72.1*	*67.2*	42.5	42.0
40-44	55.8	*59.0*	48.5	*50.3*	*96.5*	--	50.5	*52.8*
45-49	--	--	--	--	--	--	--	--
Marital Status								
Married, or in union	43.8	na	53.5	na	83.9	na	44.8	na
Unmarried, sexually active	19.3	na	24.5	na	*73.5*	na	-50.2	na
Parity								
0-1 children	18.2	28.9	23.9	41.8	69.7	--	-53.1	45.7
2-3 children	39.3	42.6	53.7	55.1	82.9	81.7	45.0	47.2
4+ children	51.5	51.1	56.6	56.6	88.7	87.5	42.7	41.9
Residence								
Urban	32.6	43.8	40.9	53.5	81.3	83.9	-0.1	44.8
Education								
Never	--	--	--	--	--	--	--	--
Primary	47.7	49.3	62.9	67.8	*97.9*	*97.6*	43.8	46.4
Secondary	34.1	45.5	43.0	53.9	82.0	82.7	-2.3	46.0
Higher	22.6	32.1	25.8	38.4	*61.1*	--	-12.3	39.3
Wealth Quintile								
Lowest	44.2	52.5	54.3	65.6	95.7	*94.0*	-83.5	46.1
Lower	35.0	46.2	49.3	60.1	76.1	*81.5*	42.4	49.3
Middle	29.6	43.8	30.6	45.6	*75.6*	--	24.2	25.3
High	27.5	42.2	29.9	45.9	*81.7*	*87.5*	-13.8	49.1
Highest	26.6	36.4	38.3	49.2	*70.8*	*73.4*	36.2	49.1
Region								
Kinshasa	32.6	43.8	40.9	53.5	81.3	83.9	-0.1	44.8

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactional amenorrhea method

2 Asked of all current/recent modern contraceptive users

3 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Quality of Care

Percentage of current modern[1] contraceptive users who would return, would refer a friend/relative, or would return and refer a friend/relative to provider, by background characteristics.

Background Characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Total	**(N = 585)**	**(N = 585)**	**(N = 585)**
All	58.6	62.5	44.6
Age Group			
15-19	45.6	45.3	25.6
20-24	61.1	59.6	46.9
25-29	64.8	66.1	50.3
30-34	67.1	62.3	48.4
35-39	49.9	72.7	44.6
40-44	*44.6*	*61.4*	*36.7*
45-49	--	--	--
Marital Status			
Married, or in union	62.8	69.4	50.7
Unmarried, sexually active	54.5	53.5	37.8
Parity			
0-1 children	56.7	51.5	39.6
2-3 children	62.5	68.6	51.4
4+ children	57.1	75.0	45.2
Residence			
Urban	58.6	62.5	44.6
Education			
Never	--	--	--
Primary	72.7	81.9	63.4
Secondary	55.0	61.8	40.4
Higher	63.3	56.4	49.8
Wealth Quintile			
Lowest	61.8	79.2	47.6
Lower	58.4	69.8	48.7
Middle	50.0	54.1	35.3
High	59.4	52.2	39.6
Highest	61.5	55.8	49.6
Region			
Kinshasa	58.6	62.5	44.6

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactional amenorrhea method

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Knowledge of Family Planning

Percentage of women age 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics.

Background Characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All Women	Married[1] Women	All Women	Married[1] Women	All Women	Married[1] Women
Total	**(2611)**	**(1114)**	**(2611)**	**(1114)**	**(2611)**	**(1114)**
All	4.1	5.1	11.8	18.9	14.9	22.5
Age Group						
15-19	4.0	*15.6*	4.0	*37.3*	7.2	*42.8*
20-24	5.6	9.8	13.8	35.3	18.3	42.7
25-29	2.7	3.6	17.0	23.0	19.0	25.5
30-34	4.8	5.3	19.7	21.4	22.6	24.8
35-39	2.8	2.9	14.4	18.7	16.4	20.4
40-44	5.2	6.8	8.8	9.4	13.0	14.8
45-49	2.7	1.8	4.8	3.7	7.0	5.5
Marital Status						
Married, or in union	5.1	na	18.9	na	22.5	na
Unmarried, sexually active	3.0	na	10.1	na	12.4	na
Parity						
0-1 children	3.3	4.6	7.3	20.1	9.7	22.1
2-3 children	4.8	4.5	19.8	20.7	23.1	23.6
4+ children	5.7	6.1	15.1	16.2	20.0	21.4
Residence						
Urban	4.1	5.1	11.8	18.9	14.9	22.5
Education						
Never	--	--	--	--	--	--
Primary	2.0	2.4	16.2	23.9	16.7	24.6
Secondary	4.9	5.8	12.6	19.1	16.4	23.4
Higher	2.1	4.1	6.7	13.2	8.2	15.8
Wealth Quintile						
Lowest	6.4	4.2	16.4	26.6	21.2	28.2
Lower	6.1	7.7	12.0	17.8	16.8	24.4
Middle	3.0	3.7	11.1	19.0	13.8	22.1
High	2.8	5.4	10.7	18.0	12.9	22.2
Highest	2.6	4.7	9.2	14.4	10.9	17.0
Region						
Kinshasa	4.1	5.1	11.8	18.9	14.9	22.5

1   *Married includes married women and those who are living with a man*

na *Not applicable*

--   *Less than 25 unweighted observations*

<i> *Italics indicate estimate was generated from 25 to 49 unweighted observations*

Response Rates

Response rates of households and individuals, by residence.

Result	Urban	Rural	Total
Household Interviews			
Households Selected	N/A	N/A	2030
Households Occupied	N/A	N/A	2009
Households Interviewed	N/A	N/A	1968
Household Response Rate [1]	0.00%	0.00%	97.96%
Interviews with Women			
Number of Eligible Women [2]	N/A	N/A	2740
Number of Eligible Women Interviewed	N/A	N/A	2611
Eligible Woman Response Rate [3]	0.00%	0.00%	95.29%

1   *Household response rate = households interviewed/households selected*

2   *Eligible women are women ages 15-49 identified in completed household interviews*

3   *Eligible response rate = eligible women interviewed/eligible women*

Sample Error Estimates

Variable	All Women				Married Women[1]			
	Mean	Std Error	CI Lower	CI Upper	Mean	Std Error	CI Lower	CI Upper
Currently using a modern method	0.245	0.014	0.218	0.274	0.296	0.024	0.251	0.346
Currently using a traditional method	0.191	0.014	0.166	0.220	0.250	0.021	0.211	0.294
Currently using any contraceptive method	0.436	0.016	0.405	0.468	0.547	0.022	0.502	0.590
Currently using DMPA-IM	0.020	0.005	0.013	0.032	0.032	0.009	0.018	0.055
Currently using DMPA-SC	0.008	0.002	0.004	0.014	0.015	0.005	0.008	0.030
Currently using condoms	0.156	0.017	0.124	0.193	0.100	0.015	0.074	0.134
Currenlty using implants	0.067	0.012	0.046	0.096	0.119	0.018	0.087	0.160
Chose method by self or jointly in past 12 months	0.869	0.026	0.809	0.913	0.898	0.022	0.846	0.934
Informed by provider about other methods	0.326	0.036	0.258	0.402	0.438	0.034	0.372	0.506
Informed by provider about side effects	0.409	0.045	0.322	0.501	0.535	0.043	0.448	0.619
Satisfied with provider: Would return and refer friend/relative to provider	0.446	0.046	0.357	0.540	0.507	0.044	0.419	0.594
Visited by health worker who talked about family planning in past 12 months	0.041	0.009	0.026	0.063	0.051	0.012	0.032	0.081

1   Married includes married women and those who are living with a man