

Background characteristics of respondents**Percent distribution of background characteristics of women age 15 to 49**

Background characteristics	Urban	Rural	Percentage of sample	Weighted number	Unweighted number
Age group					
15–19	65.0	35.0	100.0	16.0	200
20–24	65.3	34.7	100.0	14.9	186
25–29	76.8	23.2	100.0	20.8	260
30–34	74.2	25.8	100.0	18.5	232
35–39	65.2	34.8	100.0	13.1	163
40–44	55.2	44.8	100.0	11.2	139
45–49	57.5	42.5	100.0	5.6	70
Marital status					
Married	67.4	32.6	100.0	47.3	591
Living together	59.6	40.4	100.0	9.8	122
Divorced	54.1	45.9	100.0	2.6	33
Widowed	69.2	30.8	100.0	2.9	36
Never married	71.1	28.9	100.0	37.4	468
Parity					
None	73.1	26.9	100.0	40.8	510
1-2	74.9	25.1	100.0	25.8	322
3-4	64.5	35.5	100.0	18.4	230
5 or more	44.8	55.2	100.0	15.0	188
Education					
Never attended	(43.4)	(56.6)	(100.0)	1.1	13
Primary	57.8	42.2	100.0	12.8	160
Secondary	66.2	33.8	100.0	62.2	778
Higher	78.2	21.8	100.0	23.9	298
Wealth quintile					
Lowest	36.7	63.3	100.0	16.7	208
Lower	77.2	22.8	100.0	20.5	257
Middle	72.4	27.6	100.0	21.1	263
Higher	68.7	31.3	100.0	18.2	227
Highest	76.5	23.5	100.0	23.5	294
Total	67.7	32.3	100.0	100.0	1,250

Current use of contraception by background characteristics**Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics**

Background characteristic	Any method		Modern method ¹		Traditional method	
	All women	Married women	All women	Married women	All women	Married women
Number	1,250	713	1,250	713	1,250	713
Total	28.1	32.9	19.4	21.6	8.1	10.1
Age group						
15–19	9.3	(39.0)	7.3	(39.0)	2.1	(0.0)
20–24	34.4	32.0	27.0	29.6	7.4	2.4
25–29	32.1	33.1	25.4	23.5	6.7	9.6
30–34	31.2	31.0	21.9	24.1	8.3	5.6
35–39	29.7	31.6	18.0	17.6	11.7	14.0
40–44	32.2	35.8	11.3	11.9	16.6	19.1
45–49	28.4	35.7	22.3	27.9	6.1	7.8
Marital status						
Married ²	32.9	na	21.6	na	10.1	na
Unmarried, sexually active	46.5	na	32.8	na	13.7	na
Parity						
0–1	20.6	18.5	15.2	14.0	5.4	4.5
2–3	30.6	31.2	20.2	19.6	8.9	9.8
4 or more	42.8	45.9	28.0	29.5	13.4	14.8
Residence						
Urban	29.2	35.5	21.7	25.1	7.0	9.5
Rural	25.9	28.0	14.5	14.9	10.5	11.4
Education						
Never attended	(10.8)	(11.7)	(9.9)	(9.9)	(1.0)	(1.8)
Primary	20.8	22.8	16.3	18.7	3.3	2.6
Secondary	28.2	34.6	19.3	22.5	8.6	11.6
Higher	32.7	36.7	21.7	22.3	9.7	12.3
Wealth quintile						
Lowest	20.1	18.1	14.8	12.0	5.2	6.1
Lower	25.6	33.0	17.9	23.0	7.7	9.9
Middle	31.5	37.3	26.9	32.1	3.9	3.9
Higher	35.2	40.0	18.1	17.0	14.4	18.8
Highest	27.5	32.1	18.2	21.8	9.4	10.3

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Contraceptive method mix by background characteristics**Percent distribution of current contraceptive users ages 15 to 49, by method type, marital status and background characteristics**

All users										
Number = 352										
Background characteristics	Sterilization	Implants	IUD	Injectables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	0.9	7.1	1.9	9.5	17.2	6.5	25.3	3.6	27.9	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(18.5)	(24.7)	(34.7)	(0.0)	(22.0)	(100.0)
20–24	0.0	1.0	0.0	7.1	17.0	13.2	40.3	1.6	19.9	100.0
25–29	0.0	11.8	0.6	6.2	20.8	5.8	32.2	2.2	20.5	100.0
30–34	0.5	14.1	5.5	13.6	20.2	1.5	14.7	5.0	25.0	100.0
35–39	1.1	5.7	0.0	8.4	12.1	5.7	27.4	0.0	39.5	100.0
40–44	0.5	3.3	4.6	10.7	12.1	2.6	1.3	14.0	50.8	100.0
45–49	(11.1)	(0.0)	(0.0)	(25.1)	(15.2)	(0.0)	(27.1)	(0.0)	(21.4)	(100.0)
Marital status										
Married ¹	1.4	10.2	2.8	12.6	17.1	4.5	16.9	4.6	29.9	100.0
Unmarried, sexually active	0.0	0.6	0.0	1.9	18.8	4.6	43.3	1.3	29.5	100.0
Parity										
0–1	0.0	0.0	0.0	5.6	15.3	11.2	40.9	1.4	25.6	100.0
2–3	0.0	14.5	0.6	12.2	12.2	4.3	21.7	7.1	27.4	100.0
4 or more	2.5	10.0	4.7	12.0	22.5	2.9	10.8	3.8	30.7	100.0
Residence										
Urban	1.1	8.0	0.8	10.2	17.1	6.9	30.0	2.9	23.1	100.0
Rural	0.5	5.1	4.3	7.9	17.6	5.7	14.3	5.3	39.3	100.0
Education										
Never attended	(0.0)	(0.0)	(0.0)	(0.0)	(90.8)	(0.0)	(0.0)	(0.0)	(9.2)	(100.0)
Primary	0.0	8.4	0.0	34.2	25.6	2.7	7.5	6.7	14.8	100.0
Secondary	0.4	6.9	1.2	7.2	16.3	8.0	27.6	3.0	29.3	100.0
Higher	2.5	7.2	4.1	6.3	15.3	4.5	26.5	4.1	29.6	100.0
Wealth quintile										
Lowest	0.0	4.8	0.0	7.7	25.4	14.8	21.2	4.8	21.4	100.0
Lower	0.0	9.6	0.0	5.4	14.7	7.2	33.1	0.5	29.5	100.0
Middle	3.7	3.9	2.5	19.7	24.2	7.9	22.7	3.3	11.9	100.0
Higher	0.0	8.7	0.7	7.2	12.3	2.4	18.7	9.6	40.4	100.0
Highest	0.3	7.9	4.9	5.5	12.8	4.2	30.3	0.0	34.0	100.0

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Married users										
Number = 235										
Background characteristics	Sterilization	Implants	IUD	Injectables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	1.4	10.2	2.8	12.6	17.1	4.5	16.9	4.6	29.9	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(83.8)	(16.2)	(0.0)	(0.0)	(0.0)	(100.0)
20–24	0.0	0.8	0.0	19.4	21.7	17.4	33.2	4.8	2.7	100.0
25–29	0.0	18.9	1.0	5.5	13.7	4.3	26.8	1.0	28.9	100.0
30–34	0.7	18.9	7.4	18.2	24.4	1.2	6.7	5.6	16.8	100.0
35–39	1.3	5.6	0.0	8.9	11.4	6.7	21.8	0.0	44.4	100.0
40–44	0.5	3.1	4.8	10.6	11.2	1.7	1.4	14.4	52.3	100.0
45–49	(11.3)	(0.0)	(0.0)	(23.8)	(15.5)	(0.0)	(27.6)	(0.0)	(21.8)	(100.0)
Marital status										
Married ¹	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	na	na	na	na	na	na	na	na	na	na
Parity										
0–1	0.0	0.0	0.0	14.4	20.0	11.8	29.3	1.1	23.4	100.0
2–3	0.0	16.0	0.7	13.3	9.6	3.0	19.5	8.1	29.7	100.0
4 or more	2.7	10.1	4.9	11.7	20.5	3.0	11.3	3.7	32.0	100.0
Residence										
Urban	1.7	11.7	1.2	13.4	17.2	3.9	21.6	3.4	25.9	100.0
Rural	0.8	6.5	6.7	10.7	16.8	6.0	5.2	7.8	39.6	100.0
Education										
Never attended	(0.0)	(0.0)	(0.0)	(0.0)	(84.5)	(0.0)	(0.0)	(0.0)	(15.5)	(100.0)
Primary	0.0	8.6	0.0	41.7	27.8	1.5	2.3	8.2	9.9	100.0
Secondary	0.6	10.6	1.9	10.1	16.7	6.4	18.1	3.4	32.1	100.0
Higher	3.5	10.1	5.8	6.3	12.8	1.9	20.3	5.8	33.5	100.0
Wealth quintile										
Lowest	0.0	5.3	0.0	11.7	29.6	13.2	6.3	7.6	26.3	100.0
Lower	0.0	14.3	0.0	6.8	13.6	4.6	30.6	0.0	30.1	100.0
Middle	5.7	6.0	3.8	28.7	22.4	7.3	11.1	5.1	9.9	100.0
Higher	0.0	11.5	0.9	9.5	11.9	0.2	8.5	11.0	46.6	100.0
Highest	0.4	11.3	7.0	5.5	16.1	3.3	24.1	0.0	32.2	100.0


¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Current Method Mix Among All and Married Contraceptive Users (Age 15-49)



Unmet need for family planning

Percentage of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics

Background characteristics	All women			Married women		
	Number	1,250		713		Number
	For spacing	For limiting	Total	For spacing	For limiting	Total
Total	9.2	7.1	16.4	11.2	11.2	22.4
Age group						
15–19	4.9	1.3	6.2	(14.8)	(0.0)	(14.8)
20–24	13.3	1.4	14.7	18.7	0.6	19.3
25–29	13.7	4.3	18.0	15.0	7.1	22.1
30–34	14.8	2.8	17.5	18.6	3.3	22.0
35–39	3.2	13.4	16.7	3.7	15.6	19.3
40–44	3.3	24.9	28.2	3.7	27.1	30.8
45–49	2.1	13.2	15.3	2.1	16.1	18.2
Marital status						
Married ¹	11.2	11.2	22.4	na	na	na
Unmarried, sexually active	20.4	2.9	23.4	na	na	na
Parity						
0–1	8.7	0.7	9.4	12.9	0.2	13.1
2–3	13.2	4.7	17.9	14.4	4.6	19.0
4 or more	7.0	23.5	30.5	7.2	25.5	32.7
Residence						
Urban	8.6	4.7	13.3	11.2	7.3	18.5
Rural	10.6	12.2	22.9	11.3	18.6	29.9
Education						
Never attended	(5.0)	(3.9)	(8.9)	(9.2)	(7.1)	(16.2)
Primary	7.4	18.7	26.0	5.8	23.1	28.9
Secondary	10.3	5.7	16.0	14.1	9.5	23.6
Higher	7.8	4.6	12.4	8.8	7.2	16.0
Wealth quintile						
Lowest	16.2	9.9	26.1	17.0	16.0	32.9
Lower	9.5	5.5	15.0	12.4	10.2	22.6
Middle	9.4	7.2	16.6	13.8	12.3	26.1
Higher	7.5	7.9	15.4	10.6	9.7	20.4
Highest	5.4	5.8	11.2	5.3	9.3	14.6

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Need and demand for family planning

Percentage women ages 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics

Background characteristics	All women					Married women				
	1,250					713				
Number	Modern method ¹	Any method	Unmet need	Total demand	Demand satisfied by modern method	Modern method ¹	Any method	Unmet need	Total demand	Demand satisfied by modern method
Total	19.4	28.1	16.4	44.1	43.9	21.6	32.9	22.4	54.7	39.5
Age group										
15–19	7.3	9.3	6.2	15.5	46.9	(39.0)	(39.0)	(14.8)	(53.8)	(72.5)
20–24	27.0	34.4	14.7	49.1	55.0	29.6	32.0	19.3	51.3	57.6
25–29	25.4	32.1	18.0	50.0	50.7	23.5	33.1	22.1	55.2	42.7
30–34	21.9	31.2	17.5	47.7	45.8	24.1	31.0	22.0	51.6	46.6
35–39	18.0	29.7	16.7	46.4	38.8	17.6	31.6	19.3	51.0	34.5
40–44	11.3	32.2	28.2	59.1	19.2	11.9	35.8	30.8	65.0	18.3
45–49	22.3	28.4	15.3	43.7	51.0	27.9	35.7	18.2	53.8	51.8
Marital status										
Married ²	21.6	32.9	22.4	54.7	39.5	na	na	na	na	na
Unmarried, sexually active	32.8	46.5	23.4	69.9	46.9	na	na	na	na	na
Parity										
0–1	15.2	20.6	9.4	30.0	50.6	14.0	18.5	13.1	31.6	44.3
2–3	20.2	30.6	17.9	48.5	41.7	19.6	31.2	19.0	50.2	39.1
4 or more	28.0	42.8	30.5	71.9	39.0	29.5	45.9	32.7	77.0	38.3
Residence										
Urban	21.7	29.2	13.3	41.9	51.8	25.1	35.5	18.5	53.1	47.2
Rural	14.5	25.9	22.9	48.8	29.7	14.9	28.0	29.9	57.8	25.8
Education										
Never attended	(9.8)	(10.8)	(8.9)	(19.7)	(49.8)	(9.9)	(11.7)	(16.2)	(27.9)	(35.4)
Primary	16.3	20.8	26.0	45.7	35.8	18.7	22.8	28.9	50.2	37.2
Secondary	19.3	28.2	16.0	43.9	44.0	22.5	34.6	23.6	57.6	39.0
Higher	21.7	32.7	12.4	45.1	48.1	22.3	36.7	16.0	52.7	42.3
Wealth quintile										
Lowest	14.8	20.1	26.1	46.1	32.1	12.0	18.1	32.9	51.0	23.4
Lower	17.9	25.6	15.0	40.6	44.2	23.0	33.0	22.6	55.6	41.4
Middle	26.9	31.5	16.6	47.4	56.7	32.1	37.3	26.1	62.1	51.7
Higher	18.1	35.2	15.4	49.6	36.4	17.0	40.0	20.4	58.8	28.8
Highest	18.2	27.5	11.2	38.7	46.9	21.8	32.1	14.6	46.7	46.6

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Pregnancy intentions

Unintended births among women ages 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics

Background characteristics	All women with 1 or more births				Married women with 1 or more births			
	483				429			
Number	Wanted then	Wanted later	Not at all	Total	Wanted then	Wanted later	Not at all	Total
Total	63.7	19.3	17.0	100.0	67.4	18.9	13.7	100.0
Age group								
15–19	(22.9)	(15.2)	(61.9)	(100.0)	(64.1)	(35.9)	(0.0)	(100.0)
20–24	56.7	22.9	20.4	100.0	70.5	23.7	5.8	100.0
25–29	61.5	29.0	9.5	100.0	62.6	28.3	9.1	100.0
30–34	69.5	21.2	9.4	100.0	70.3	20.7	9.0	100.0
35–39	73.1	7.5	19.4	100.0	73.8	7.5	18.7	100.0
40–44	58.1	6.7	35.2	100.0	60.2	6.1	33.7	100.0
45–49	(28.6)	(0.0)	(71.4)	(100.0)	(31.7)	(0.0)	(68.3)	(100.0)
Marital status								
Married ¹	67.4	18.9	13.7	100.0	na	na	na	na
Unmarried, sexually active	31.1	24.1	44.7	100.0	na	na	na	na
Parity								
0–1	69.9	17.9	12.2	100.0	81.6	16.6	1.8	100.0
2–3	67.2	26.8	6.0	100.0	68.4	27.3	4.3	100.0
4 or more	54.8	13.4	31.7	100.0	56.0	12.9	31.1	100.0
Residence								
Urban	68.1	18.9	13.1	100.0	70.4	18.4	11.2	100.0
Rural	55.1	20.2	24.7	100.0	60.9	20.0	19.1	100.0
Education								
Never attended	(60.6)	(20.3)	(19.1)	(100.0)	(82.1)	(10.8)	(7.1)	(100.0)
Primary	50.7	16.6	32.6	100.0	49.5	18.3	32.2	100.0
Secondary	62.0	20.1	18.0	100.0	67.1	20.0	13.0	100.0
Higher	75.1	19.0	5.9	100.0	77.3	17.1	5.6	100.0
Wealth quintile								
Lowest	50.1	24.3	25.6	100.0	55.7	25.6	18.6	100.0
Lower	66.9	15.2	17.9	100.0	67.7	15.1	17.2	100.0
Middle	58.1	26.3	15.5	100.0	59.7	25.0	15.3	100.0
Higher	63.1	16.3	20.6	100.0	68.4	17.1	14.5	100.0
Highest	78.9	15.3	5.8	100.0	79.8	14.4	5.8	100.0

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Contraceptive choice

Percent distribution of women ages 15 to 49 who are currently using modern contraception¹ reporting on who decided on the contraceptive method, by background characteristics

Background characteristics	Self	Jointly with partner or provider	Provider alone/ partner alone/other	Total
Number	135	182	34	351
Total	38.5	51.8	9.7	100.0
Age group				
15–19	(33.5)	(28.6)	(37.9)	(100.0)
20–24	37.4	48.7	13.9	100.0
25–29	36.6	55.0	8.4	100.0
30–34	50.7	45.1	4.2	100.0
35–39	36.6	53.0	10.4	100.0
40–44	28.9	70.1	0.9	100.0
45–49	(37.4)	(49.8)	(12.8)	(100.0)
Marital status				
Married ²	33.1	60.6	6.3	100.0
Unmarried, sexually active	39.5	42.4	18.1	100.0
Parity				
0–1	43.5	43.0	13.5	100.0
2–3	28.5	59.1	12.4	100.0
4 or more	39.6	56.6	3.8	100.0
Residence				
Urban	41.7	47.0	11.3	100.0
Rural	31.0	63.1	5.9	100.0
Education				
Never attended	(100.0)	(0.0)	(0.0)	(100.0)
Primary	35.8	58.9	5.3	100.0
Secondary	42.1	46.2	11.7	100.0
Higher	30.5	62.7	6.7	100.0
Wealth quintile				
Lowest	55.7	28.4	15.9	100.0
Lower	38.9	44.3	16.8	100.0
Middle	38.7	51.7	9.7	100.0
Higher	31.2	64.4	4.4	100.0
Highest	36.5	57.6	5.9	100.0

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Payment for family planning

Percentage of women ages 15 to 49 currently using a modern contraceptive method¹ who paid fees for services, by marital status and background characteristics

Background characteristics	All women who used in past 12 months	All married women who used in past 12 month
Number	352	235
Total	35.8	36.5
Age group		
15–19	(52.7)	(100.0)
20–24	42.0	52.8
25–29	44.4	46.1
30–34	38.7	46.2
35–39	23.7	25.7
40–44	14.2	12.9
45–49	(32.3)	(32.9)
Marital status		
Married ²	36.5	na
Unmarried, sexually active	30.5	na
Parity		
0–1	38.6	52.5
2–3	41.5	42.9
4 or more	29.3	27.7
Residence		
Urban	41.5	41.5
Rural	22.6	24.2
Education		
Never attended	(9.2)	(15.5)
Primary	44.1	49.6
Secondary	36.6	39.1
Higher	31.7	26.4
Wealth quintile		
Lowest	41.5	36.6
Lower	32.0	32.2
Middle	45.9	50.7
Higher	25.5	26.9
Highest	35.9	36.6

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Method Information Index Components

Percentage of current modern¹ contraceptive users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics²

Background characteristics	Informed about other methods		Informed about side effects		Told what to do if experienced side effects	
	All users	Married users	All users	Married users	All users	Married users
Number	352	235	250	156	71	50
Total	41.9	52.8	28.5	32.2	81.3	84.4
Age group						
15–19	(25.7)	(0.0)	(26.5)	(0.0)	(86.3)	(0.0)
20–24	14.9	24.8	9.8	(7.4)	(30.3)	(71.0)
25–29	40.3	47.0	35.7	44.4	(74.9)	(68.8)
30–34	39.0	46.2	31.7	38.4	(94.0)	(95.1)
35–39	64.2	71.0	29.0	15.7	(84.0)	(74.1)
40–44	63.5	64.9	(32.5)	(31.7)	(86.2)	(88.9)
45–49	(58.0)	(59.0)	(49.4)	(50.5)	(100.0)	(100.0)
Marital status						
Married ³	52.8	na	32.2	na	84.4	na
Unmarried, sexually active	20.5	na	12.5	na	(73.1)	na
Parity						
0–1	22.6	34.3	17.4	16.1	(63.0)	(36.8)
2–3	39.2	40.3	25.5	23.2	(83.3)	(88.1)
4 or more	64.3	65.9	44.3	43.1	89.7	89.9
Residence						
Urban	40.7	48.9	27.6	32.5	84.1	86.2
Rural	44.7	62.3	31.1	31.0	74.0	77.8
Education						
Never attended	(81.6)	(68.9)	(0.0)	(0.0)	na	na
Primary	53.9	45.0	44.6	(37.1)	(64.8)	(61.7)
Secondary	34.1	46.2	24.4	26.1	78.8	82.1
Higher	54.7	68.7	32.2	42.9	(94.9)	(97.6)
Wealth quintile						
Lowest	37.8	42.1	30.8	32.3	(64.4)	(68.8)
Lower	29.2	29.8	33.2	46.3	(78.6)	(83.1)
Middle	45.2	57.5	25.4	23.1	(69.8)	(71.0)
Higher	41.4	51.4	10.5	(13.0)	(81.1)	(73.6)
Highest	51.4	71.2	42.1	(44.8)	(100.0)	(100.0)

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Informed about other methods" asked of all current contraceptive users, "informed about side effects" asked of current modern contraceptive users, and "told what to do if experienced side effects" asked of current modern contraceptive users who indicated they were told about side effects.

³Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Perceived quality of care for family planning

Percentage of current modern¹ contraceptive users who would return, would refer a friend/relative, or would return and refer friend/relative to provider, by background characteristics

Background characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Number	249	249	249
Total	68.6	64.7	55.8
Age group			
15–19	(43.4)	(22.4)	(15.4)
20–24	65.8	55.1	51.6
25–29	72.2	71.6	60.8
30–34	68.5	79.7	60.1
35–39	81.2	63.1	61.7
40–44	(48.7)	(45.3)	(43.8)
45–49	(83.9)	(82.5)	(75.2)
Marital status			
Married ²	74.7	73.3	63.6
Unmarried, sexually active	53.1	49.4	38.9
Parity			
0–1	62.0	49.1	45.7
2–3	76.8	79.8	66.3
4 or more	71.8	75.0	62.0
Residence			
Urban	70.6	66.7	58.2
Rural	62.7	58.9	48.7
Education			
Never attended	(100.0)	(100.0)	(100.0)
Primary	67.6	61.5	50.7
Secondary	69.0	62.4	55.1
Higher	67.4	70.7	58.8
Wealth quintile			
Lowest	74.1	76.7	64.6
Lower	85.5	73.2	70.6
Middle	66.7	67.3	59.1
Higher	53.3	42.8	38.1
Highest	65.7	65.1	48.1

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Knowledge of family planning

Percentage of women ages 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics

Background characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All women	Married women ¹	All women	Married women ¹	All women	Married women ¹
Number	1,250	713	1,250	713	1,250	713
Total	8.5	11.0	16.4	24.9	20.9	30.2
Age group						
15–19	4.8	(0.0)	1.1	(0.0)	5.0	(0.0)
20–24	4.6	11.0	9.2	19.0	10.2	20.3
25–29	9.2	8.4	19.6	26.7	25.3	31.9
30–34	11.7	14.8	28.1	34.8	34.4	42.5
35–39	9.3	10.5	24.6	30.0	27.2	32.6
40–44	13.3	14.6	16.0	14.5	24.4	24.2
45–49	3.5	0.6	9.4	9.3	12.4	9.9
Marital status						
Married ¹	11.0	na	24.9	na	30.2	na
Unmarried, sexually active	3.7	na	5.0	na	8.0	na
Parity						
0–1	7.0	11.3	7.9	18.2	12.4	24.6
2–3	11.1	11.9	28.1	30.3	33.2	35.7
4 or more	9.3	9.9	24.8	25.5	29.1	29.9
Residence						
Urban	7.3	10.4	17.2	27.1	20.8	32.1
Rural	10.9	12.1	14.7	20.6	21.3	26.5
Education						
Never attended	(3.0)	(0.0)	(7.0)	(4.1)	(7.0)	(4.1)
Primary	7.3	7.1	12.4	13.1	17.5	18.1
Secondary	7.5	10.9	13.7	23.0	17.7	28.3
Higher	11.7	14.0	25.7	37.1	32.0	43.1
Wealth quintile						
Lowest	7.8	9.7	11.4	13.4	15.8	18.0
Lower	9.7	12.5	12.7	20.2	18.0	25.8
Middle	8.3	10.8	13.9	20.6	18.5	27.1
Higher	12.1	14.2	21.6	29.1	30.3	39.3
Highest	5.1	7.8	21.2	35.1	22.0	35.7

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

List of indicators for sampling errors, PMA2016/Rivers

Variable	Estimation	Base population
Currently using a modern method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using a traditional method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using any contraceptive method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using injectables	Proportion	All women ages 15–49 Married women ages 15–49
Currently using male condoms	Proportion	All women ages 15–49 Married women ages 15–49
Currently using implants	Proportion	All women ages 15–49 Married women ages 15–49
Chose method by self or jointly in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Paid fees for family planning services in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about other methods	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about side effects	Proportion	All women ages 15–49 Married women ages 15–49
Satisfied with provider: Would return and refer friend/relative to provider	Proportion	All women ages 15–49 Married women ages 15–49
Visited by health worker who talked about family planning information in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49

Characteristics of service delivery points

Number and percentage of service delivery points (SDPs) by background characteristics

Background characteristics	Number	Percentage
Total	111	100.0
SDP type		
Hospital	17	15.3
Health center ¹	37	33.3
Pharmacy ²	57	51.4
Sector		
Public	51	45.9
Private	60	54.1
Offers family planning		
No	17	15.3
Yes	94	84.7

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

Charging fees for family planning

Percentage of service delivery points (SDPs) that charge fees for family planning services, by sector and background characteristics, among SDPs offering family planning

Background characteristic	Sector	
	Public	Private
Number of SDPs	47	47
Total	34.0	51.1
SDP type		
Hospital	(58.3)	(100.0)
Health center ¹	25.7	(0.0)
Pharmacy ²	na	52.3

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive choice: Availability of at least 3 or at least 5 modern contraceptive methods

Percentage of public and private service delivery points (SDPs) offering at least 3 or 5 modern contraceptive methods¹, by sector and background characteristics, among all SDPs

Facility characteristic	3 or more methods		5 or more methods	
	Public	Private	Public	Private
Number of SDPs	51	60	51	60
Total	82.4	26.7	74.5	3.3
Facility type				
Hospital	(68.8)	(100.0)	(62.5)	(0.0)
Health Center ²	88.6	(50.0)	80.0	(0.0)
Pharmacy ³	na	24.6	na	3.5

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive choice: Availability of modern contraception, by method

Percentage of service delivery points (SDPs) offering different types of contraceptive methods¹, by sector and method, among all SDPs

Public SDPs							
Contraceptive method							
SDP characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number = 51							
Total	88.2	84.3	70.6	82.4	88.2	13.7	25.5
Facility type							
Hospital	(68.8)	(56.3)	(50.0)	(62.5)	(68.8)	(43.8)	(25.0)
Health center ²	97.1	97.1	80.0	91.4	97.1	0.0	25.7
Pharmacy ³	na	na	na	na	na	na	na

Private SDPs							
Contraceptive method							
SDP characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number = 60							
Total	55.0	23.3	1.7	0.0	71.7	1.7	35.0
Facility type							
Hospital	(100.0)	(100.0)	(100.0)	(0.0)	(0.0)	(100.0)	(0.0)
Health center ²	(50.0)	(100.0)	(0.0)	(0.0)	(50.0)	(0.0)	(100.0)
Pharmacy ³	54.4	19.3	0.0	0.0	73.7	0.0	33.3

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

(XX) = Estimate generated from less than 25 unweighted observations

(XX) = Estimate generated from less than 5 unweighted observations

Contraceptive stock-outs, by method

Percentage of service delivery points (SDPs) stocked out of specific modern contraceptives¹ at time of survey or stocked out in the last 3 months, by sector and background characteristics, among SDPs that offer each method

Public SDPs						
Contraceptive method						
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	45	43	36	42	45	13
Total	13.3	9.3	8.3	4.8	11.1	(46.2)
SDP type						
Hospital	(18.2)	(0.0)	(12.5)	(0.0)	(9.1)	(50.0)
Health center ²	11.8	11.8	7.1	6.3	11.8	(44.4)
Pharmacy ³	na	na	na	na	na	na

Private SDPs						
Contraceptive method						
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	33	14	1	0	43	21
Total	27.3	(57.1)	(0.0)	na	9.3	(23.8)
SDP type						
Hospital	(0.0)	(0.0)	(0.0)	na	na	na
Health center ²	(0.0)	(100.0)	na	na	(0.0)	(0.0)
Pharmacy ³	29.0	(54.5)	na	na	9.5	(26.3)

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Number of family planning visits, by method

Total number of visits and new clients for contraceptive services, by method type and background characteristics, among advanced¹ service delivery points (SDPs) that offer each method

	Contraceptive Methods													
	Sterilization		Injectable		IUD		Implants		Male condom		Pills		Emergency Contraception	
	Total women	Total men	Total	New	Total	New	Total	New	Total	New	Total	New	Total	New
Number of SDPs	8	2	46		37		42		46		47		15	
Total	2	0	1,246	495	77	42	313	229	2,535	1,418	193	95	11	11
Background characteristic														
SDP type														
Hospital	2	0	66	15	38	10	59	38	89	30	21	14	7	7
Health center ¹	na	na	1,180	480	39	32	254	191	2,446	1,388	172	81	4	4
Sector														
Public	0	0	1,096	485	77	42	313	229	2,535	1,418	191	95	11	11
Private	2	na	150	10	0	0	0	0	0	0	2	0	0	0

¹Includes Health Post, Health Clinic, and Maternity Center