# **Background characteristics of respondents**

## Percent distribution of background characteristics of women age 15 to 49

Background characteristics	Urban	Rural	Percentage of sample	Weighted number	Unweighted number
Age group					
15–19	21.1	78.9	100.0	23.4	199
20–24	23.8	76.2	100.0	18.0	153
25–29	15.0	85.0	100.0	18.8	160
30–34	23.2	76.8	100.0	15.1	128
35–39	15.0	85.0	100.0	10.7	91
40–44	9.9	90.1	100.0	9.6	82
45–49	20.3	79.7	100.0	4.4	37
Marital status					
Married	13.6	86.4	100.0	68.2	580
Living together	(63.3)	(36.7)	(100.0)	1.4	12
Divorced	0.0	100.0	100.0	4.5	39
Widowed	(31.6)	(68.4)	(100.0)	1.9	17
Never married	34.6	65.4	100.0	23.8	203
Parity					
None	30.8	69.2	100.0	28.8	245
1-2	17.4	82.6	100.0	28.3	241
3-4	14.8	85.2	100.0	18.6	158
5 or more	10.1	89.9	100.0	24.2	206
Education					
Never attended	6.1	93.9	100.0	38.0	323
Primary	10.5	89.5	100.0	25.6	218
Secondary	33.5	66.5	100.0	31.5	268
Higher	69.9	30.1	100.0	4.8	41
Wealth quintile					
Lowest	100.0	0.0	100.0	17.1	146
Lower	0.7	99.3	100.0	19.8	168
Middle	10.9	89.1	100.0	20.1	171
Higher	16.9	83.1	100.0	19.9	169
Highest	57.4	42.6	100.0	23.2	197
Total	19.0	81.0	100.0	100.0	850

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

#### Current use of contraception by background characteristics

## Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics

	Any m	ethod	Modern	method <sup>1</sup>	Tradition	al method
Background characteristic	All women	Married <sup>2</sup> women	All women	Married <sup>2</sup> women	All women	Married <sup>2</sup> women
Number	850	592	850	592	850	592
Total	12.9	12.5	9.9	9.6	2.9	2.9
Age group						
15–19	9.8	6.2	7.1	4.5	2.8	1.7
20–24	9.3	4.2	7.0	2.1	2.3	2.1
25–29	13.2	11.4	9.8	8.8	3.4	2.6
30–34	16.3	16.3	14.1	13.8	2.1	2.5
35–39	17.0	18.8	13.6	14.6	3.4	4.2
40–44	16.2	18.9	11.6	13.5	4.6	5.4
45–49	13.2	16.1	11.1	13.6	2.1	2.5
Marital status						
Married <sup>2</sup>	12.5	na	9.6	na	2.9	na
Unmarried, sexually active	38.4	na	28.9	na	9.5	na
Parity						
0–1	10.4	5.4	8.0	4.6	2.4	0.8
2–3	11.1	10.4	8.1	7.1	2.9	3.3
4 or more	17.6	18.4	13.9	14.4	3.6	3.9
Residence						
Urban	22.1	23.8	19.6	23.8	2.5	0.0
Rural	10.7	10.6	7.7	7.2	3.0	3.4
Education						
Never attended	6.0	6.0	5.0	5.2	1.0	0.8
Primary	12.9	15.6	9.4	11.5	3.4	4.2
Secondary	18.6	17.9	14.2	13.5	4.4	4.4
Higher	29.6	(39.8)	23.8	(28.7)	5.7	(11.1)
Wealth quintile		, ,				
Lowest	7.6	8.0	5.7	6.1	2.0	1.9
Lower	14.2	12.4	8.2	6.1	5.9	6.3
Middle	10.0	9.1	6.9	5.1	3.1	3.9
Higher	10.0	13.1	8.4	10.8	1.6	2.3
Highest	20.5	20.1	18.5	20.1	2.0	0.0

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

<sup>&</sup>lt;sup>2</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

# Contraceptive method mix by background characteristic

## Percent distribution of current contraceptive users ages 15 to 49, by method type and background characteristics

	All users									
					Nu	mber = 109				
Background characteristics	Steriliz- ation	Implants	IUD	Inject- ables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	2.1	3.0	0.0	26.1	18.2	2.1	23.6	8.3	16.7	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(28.9)	(0.0)	(43.0)	(6.0)	(22.1)	(100.0)
20–24	(0.0)	(0.0)	(0.0)	(15.7)	(15.0)	(0.0)	(44.3)	(11.0)	(14.0)	(100.0)
25–29	(0.0)	(12.6)	(0.0)	(9.4)	(31.0)	(5.6)	(10.2)	(11.1)	(20.1)	(100.0)
30–34	(7.5)	(0.0)	(0.0)	(35.3)	(16.5)	(5.6)	(22.0)	(5.7)	(7.5)	(100.0)
35–39	(0.0)	(0.0)	(0.0)	(61.3)	(8.6)	(0.0)	(10.1)	(0.0)	(20.1)	(100.0)
40–44	(5.7)	(4.6)	(0.0)	(41.9)	(5.9)	(0.0)	(8.9)	(16.4)	(16.7)	(100.0)
45–49	(0.0)	(0.0)	(0.0)	(40.2)	(0.0)	(0.0)	31.7	(12.3)	(15.8)	(100.0)
Marital status										
Married	3.1	4.4	0.0	35.4	19.0	3.2	8.3	12.2	14.4	100.0
Unmarried, sexually active	(0.0)	(0.0)	(0.0)	(2.8)	(14.1)	(0.0)	(58.4)	(0.0)	(24.7)	(100.0)
Parity										
0–1	0.0	0.0	0.0	2.1	23.8	3.1	48.2	3.1	19.8	100.0
2–3	(0.0)	(14.2)	(0.0)	(21.3)	(22.9)	(0.0)	(9.8)	(22.2)	(9.6)	(100.0)
4 or more	4.8	0.0	0.0	47.5	11.4	2.4	10.5	5.7	17.5	100.0
Residence										
Urban	0.0	7.5	0.0	25.2	32.6	0.0	23.5	0.0	11.2	100.0
Rural	3.1	8.0	0.0	26.6	11.2	3.2	23.6	12.2	19.3	100.0
Education	(2.0)	(0.0)	(0.0)	(40.0)	(15.5)	(0.0)	(1.1.0)	(0.4)	(O. O.)	(100.0)
Never attended	(3.9)	(0.0)	(0.0)	(48.2)	(17.7)	(0.0)	(14.0)	(8.1)	(8.2)	(100.0)
Primary Secondary	5.6 0.0	0.0 2.7	$0.0 \\ 0.0$	35.9 16.3	15.1 16.5	4.2 2.4	12.5 34.8	5.6 5.9	21.2 21.5	100.0 100.0
Higher	(0.0)	(15.9)	(0.0)	(9.1)	(32.7)	(0.0)	(18.1)	(24.3)	(0.0)	(100.0)
Wealth quintile	(0.0)	(13.9)	(0.0)	(9.1)	(32.1)	(0.0)	(10.1)	(24.3)	(0.0)	(100.0)
Lowest	(14.0)	(0.0)	(0.0)	(26.6)	(23.3)	(0.0)	(10.2)	(14.0)	(11.8)	(100.0)
Lower	(0.0)	(0.0)	(0.0)	(8.8)	(6.7)	(0.0)	(42.6)	(6.6)	(35.3)	(100.0)
Middle	(0.0)	(0.0)	(0.0)	(32.5)	(0.7) $(11.3)$	(0.0)	(18.2)	(20.6)	(17.4)	(100.0) $(100.0)$
Higher	(4.5)	(0.0)	(0.0)	(38.9)	(12.5)	(13.9)	(10.5)	(10.5)	(9.2)	(100.0)
Highest	0.0	8.1	0.0	28.1	28.8	0.0	23.7	1.5	9.8	100.0

<sup>&</sup>lt;sup>1</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>( ) =</sup> Estimate generated from less than 5 unweighted observations

	Married users									
						N = 74				
Background characteristics	Steriliz- ation	Implants	IUD	Inject- ables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	3.1	4.4	0.0	35.4	19.0	3.2	8.3	12.2	14.4	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(72.5)	(0.0)	(0.0)	(27.5)	(0.0)	(100.0)
20–24	(0.0)	(0.0)	(0.0)	(32.3)	(17.9)	(0.0)	(0.0)	(34.9)	(14.9)	(100.0)
25–29	(0.0)	(17.6)	(0.0)	(13.2)	(30.5)	(7.8)	(0.0)	(15.6)	(15.3)	(100.0)
30–34	(8.6)	(0.0)	(0.0)	(40.5)	(19.0)	(6.5)	(10.4)	(6.5)	(8.6)	(100.0)
35–39	(0.0)	(0.0)	(0.0)	(56.9)	(9.5)	(0.0)	(11.2)	(0.0)	(22.4)	(100.0)
40–44	(5.7)	(4.6)	(0.0)	(41.9)	(5.9)	(0.0)	(8.9)	(16.4)	(16.7)	(100.0)
45–49	(0.0)	(0.0)	(0.0)	(40.2)	(0.0)	(0.0)	(31.7)	(12.3)	(15.8)	(100.0)
Marital status	` ,	` /		` '	` '	,	,	,	,	
Married <sup>1</sup>	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	na	na	na	na	na	na	na	na	na	na
Parity										
0–1	(0.0)	(0.0)	(0.0)	(9.5)	(62.4)	(14.0)	(0.0)	(14.0)	(0.0)	(100.0)
2–3	(0.0)	(17.1)	(0.0)	(21.5)	(17.3)	(0.0)	(5.8)	(26.7)	(11.6)	(100.0)
4 or more	5.0	0.0	0.0	45.8	11.8	2.5	10.9	5.9	18.1	100.0
Residence										
Urban	(0.0)	(13.0)	(0.0)	(43.8)	(37.8)	(0.0)	(5.4)	(0.0)	(0.0)	(100.0)
Rural	4.3	1.1	0.0	32.2	11.8	4.4	9.5	16.8	19.9	100.0
Education										
Never attended	(4.6)	(0.0)	(0.0)	(51.4)	(20.6)	(0.0)	(9.4)	(9.4)	(4.7)	(100.0)
Primary	6.0	0.0	0.0	32.9	16.4	4.5	13.6	6.0	20.5	100.0
Secondary	0.0	5.7	0.0	35.0	21.8	5.1	0.0	12.7	19.7	100.0
Higher	(0.0)	(22.9)	(0.0)	(13.1)	(15.8)	(0.0)	(13.1)	(35.1)	(0.0)	(100.0)
Wealth quintile										
Lowest	(16.7)	(0.0)	(0.0)	(31.7)	(27.7)	(0.0)	(0.0)	(16.7)	(7.2)	(100.0)
Lower	(0.0)	(0.0)	(0.0)	(13.6)	(10.4)	(0.0)	(25.3)	(10.1)	(40.5)	(100.0)
Middle	(0.0)	(0.0)	(0.0)	(45.5)	(0.0)	(0.0)	(0.0)	(33.7)	(20.8)	(100.0)
Higher	(4.9)	(0.0)	(0.0)	(37.3)	(13.6)	(15.2)	(7.6)	(11.5)	(10.0)	(100.0)
Highest	(0.0)	(14.0)	(0.0)	(45.6)	(33.2)	(0.0)	(4.7)	(2.6)	(0.0)	(100.0)


<sup>&</sup>lt;sup>1</sup>Married includes married women and those who are living with a man

na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>() =</sup> Estimate generated from less than 5 unweighted observations

# Current Method Mix Among **All and Married** Contraceptive Users (Age 15-49)



# **Unmet need for family planning**

Percentage of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics

Background characteristics		All women			Married women	
Number		850			1,300	
1 (unitoci	For spacing	For limiting	Total	For spacing	For limiting	Total
Total	21.3	5.8	27.1	25.8	7.3	33.2
Age group						
15–19	20.6	1.2	21.8	37.2	1.8	39.0
20–24	28.1	1.5	29.7	33.8	1.0	34.8
25–29	22.4	3.7	26.1	25.0	4.0	29.0
30–34	25.8	7.2	33.0	29.1	8.2	37.4
35–39	14.6	17.4	32.1	16.4	18.1	34.5
40–44	17.8	8.3	26.1	19.2	9.7	28.8
45–49	2.1	17.2	19.3	2.5	21.0	23.6
Marital status		12	15.10	2.0	21.0	20.0
Married <sup>1</sup>	25.8	7.3	33.2	na	na	na
Unmarried, sexually	38.7	6.1	44.8	na	na	na
active	20.7	0.1	11.0	ina ina	iiu	iiu
Parity						
0–1	15.6	1.0	16.6	19.5	0.8	20.3
2–3	28.7	3.6	32.3	32.1	3.4	35.5
4 or more	23.6	13.8	37.4	25.2	14.1	39.3
Residence	23.0	13.0	37.4	25.2	17.1	37.3
Urban	17.6	2.9	20.5	22.5	4.0	26.5
Rural	22.2	6.4	28.6	26.4	7.9	34.3
Education	22.2	0.1	20.0	20.4	1.5	54.5
Never attended	25.6	7.7	33.3	28.5	8.3	36.8
Primary	21.1	7.6	28.7	24.8	8.6	33.4
Secondary	17.6	2.8	20.4	23.6	4.9	28.4
Higher	13.5	0.0	13.5	(13.7)	(0.0)	(13.7)
Wealth quintile	13.3	0.0	13.3	(13.1)	(0.0)	(13.7)
Lowest	23.9	10.3	34.2	22.1	10.7	32.7
Lower	22.5	8.2	30.6	27.6	10.6	38.2
Middle	26.9	4.5	31.4	35.2	4.5	39.8
Higher	19.0	4.3	23.3	24.2	6.1	30.3
Highest	15.7	2.7	18.4	20.1	4.5	24.6
Highest	15.1	2.1	10.7	20.1	7.3	44.0

<sup>&</sup>lt;sup>1</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

#### Need and demand for family planning

Percentage women ages 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics

Background characteristics			All wome	en		Married women				
Number			850					592		
	Modern method <sup>1</sup>	Any method	Unmet need	Total demand	Demand satisfied by modern method	Modern method <sup>2</sup>	Any method	Unmet need	Total demand	Demand satisfied by modern method
Total	9.9	12.9	27.1	40.0	24.9	9.6	12.5	33.2	45.7	21.0
Age group 15–19 20–24 25–29 30–34	7.1 7.0 9.8 14.1	9.8 9.3 13.2 16.3	21.8 29.7 26.1 33.0	31.6 39.0 39.3 49.2	22.4 17.9 25.0 28.7	4.5 2.1 8.8 13.8	6.2 4.2 11.4 16.3	39.0 34.8 29.0 37.4	45.3 39.1 40.4 53.7	10.0 5.5 21.7 25.8
35–39 40–44 45–49	13.6 11.6 11.1	17.0 16.2 13.2	32.1 26.1 19.3	49.1 42.2 32.5	27.6 27.4 34.2	14.6 13.5 13.6	18.8 18.9 16.1	34.5 28.8 23.6	53.4 47.7 39.7	27.4 28.3 34.2
Marital status										
Married <sup>2</sup> Unmarried, sexually active	9.6 28.9	12.5 38.4	33.2 44.8	45.7 83.2	21.0 34.7	na na	na na	na na	na na	na na
Parity										
0–1 2–3 4 or more	8.0 8.1 13.9	10.4 11.1 17.6	16.6 32.3 37.4	27.0 43.3 54.9	29.8 18.7 25.3	4.6 7.1 14.4	5.4 10.4 18.4	20.3 35.5 39.3	25.7 45.9 57.7	18.1 15.4 25.0
Residence										
Urban Rural	19.6 7.7	22.1 10.7	20.5 28.6	42.6 39.3	46.0 19.5	23.8 7.2	23.8 10.6	26.5 34.3	50.2 44.9	47.3 16.0
Education Never attended	5.0	6.0	33.3	39.3	12.7	5.2	6.0	36.8	42.8	12.1
Primary Secondary Higher	9.4 14.2 23.8	12.9 18.6 29.6	28.7 20.4 13.5	41.6 39.0 43.0	22.7 36.4 55.4	11.5 13.5 (28.7)	15.6 17.9 (39.8)	33.4 28.4 (13.7)	49.0 46.4 (53.5)	23.4 29.1 (53.6)
Wealth quintile										
Lowest Lower Middle	5.7 8.2 6.9	7.6 14.2 10.0	34.2 30.6 31.4	41.8 44.8 41.4	13.5 18.4 16.7	6.1 6.1 5.1	8.0 12.4 9.1	32.7 38.2 39.8	40.7 50.6 48.8	14.9 12.1 10.5
Higher Highest	8.4 18.5	10.0 20.5	23.3 18.4	33.3 38.9	25.2 47.6	10.8 20.1	13.1 20.1	30.3 24.6	43.4 44.7	24.8 44.9

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

<sup>&</sup>lt;sup>2</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

#### **Pregnancy intentions**

Unintended births among women ages 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics

Background characteristics	All wo	men with	1 or more	births	Married women with 1 or more births			
Number		47	<b>72</b>			47	72	
	Wanted then	Wanted later	Not at all	Total	Wanted then	Wanted later	Not at all	Total
Total	62.2	32.3	5.6	100.0	63.1	31.9	5.1	100.0
Age group								
15–19	74.4	21.9	3.7	100.0	78.3	19.6	2.1	100.0
20–24	69.1	27.2	3.7	100.0	72.2	25.8	2.0	100.0
25–29	65.7	32.4	1.8	100.0	68.2	30.2	1.5	100.0
30–34	57.7	34.6	7.7	100.0	57.9	35.2	7.0	100.0
35–39	55.6	37.2	7.2	100.0	49.7	41.8	8.4	100.0
40–44	33.4	45.5	21.1	100.0	32.0	46.5	21.5	100.0
45–49	(17.8)	(67.5)	(14.7)	(100.0)	(17.8)	(67.5)	(14.7)	(100.0)
Marital status								
Married <sup>1</sup>	63.1	31.9	5.1	100.0	na	na	na	na
Unmarried, sexually active	(31.5)	(53.3)	(15.2)	(100.0)	na	na	na	na
Parity								
0-1	78.3	17.7	4.0	100.0	85.0	15.0	0.0	100.0
2–3	72.8	25.8	1.4	100.0	75.3	23.2	1.5	100.0
4 or more	42.6	47.1	10.3	100.0	40.8	48.3	10.8	100.0
Residence								
Urban	73.1	19.3	7.6	100.0	78.9	14.8	6.2	100.0
Rural	60.6	34.2	5.3	100.0	60.6	34.5	4.9	100.0
Education								
Never attended	62.7	31.9	5.4	100.0	62.8	31.4	5.8	100.0
Primary	58.7	35.4	5.9	100.0	57.4	37.6	5.0	100.0
Secondary	64.2	30.6	5.2	100.0	68.6	27.1	4.2	100.0
Higher	(68.3)	(21.9)	(9.8)	(100.0)	(75.8)	(24.2)	(0.0)	(100.0)
Wealth quintile								
Lowest	63.8	25.7	10.5	100.0	64.1	25.5	10.3	100.0
Lower	47.0	49.7	3.3	100.0	46.6	50.5	3.0	100.0
Middle	65.1	28.3	6.7	100.0	67.2	27.9	4.8	100.0
Higher	68.4	28.7	2.8	100.0	69.4	27.4	3.2	100.0
Highest	65.5	29.7	4.8	100.0	66.4	29.6	4.0	100.0

<sup>&</sup>lt;sup>1</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

### **Contraceptive choice**

Percent distribution of women ages 15 to 49 who are currently using modern contraception reporting on who decided on the contraceptive method, by background characteristics

Background characteristics	Self	Jointly with partner or provider	Provider alone/ partner alone/other	Total
Number	20	77	12	109
Total	18.2	70.5	11.2	100.0
Age group				
15–19	(12.3)	(70.7)	(17.0)	(100.0)
20–24	(19.1)	(70.0)	(11.0)	(100.0)
25–29	(18.5)	(68.5)	(12.9)	(100.0)
30–34	(24.2)	(65.0)	(10.8)	(100.0)
35–39	(20.9)	(70.5)	(8.6)	(100.0)
40–44	(14.8)	(76.9)	(8.3)	(100.0)
45–49	(13.2)	(86.8)	(0.0)	(100.0)
Marital status		, ,		, ,
Married <sup>2</sup>	18.4	70.7	10.9	100.0
Unmarried, sexually active	(22.8)	(65.8)	(11.3)	(100.0)
Parity	(==)	()	()	()
0-1	17.5	69.7	12.8	100.0
2–3	(23.0)	(65.1)	(11.9)	(100.0)
4 or more	16.5	73.8	9.7	100.0
Residence				
Urban	14.6	79.2	6.2	100.0
Rural	19.9	66.4	13.7	100.0
Education				
Never attended	(29.3)	(52.2)	(18.4)	(100.0)
Primary	14.9	79.4	5.6	100.0
Secondary	15.2	72.7	12.2	100.0
Higher Wealth quintile	(20.6)	(70.4)	(9.1)	(100.0)
Lowest	(32.1)	(67.9)	(0.0)	(100.0)
Lower	(5.8)	(67.9)	(26.3)	(100.0)
Middle	(29.5)	(63.6)	(6.9)	(100.0)
Higher	(24.2)	(75.8)	(0.0)	(100.0)
	, ,			` /
Highest	14.4	73.6	12.0	100.0

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

<sup>&</sup>lt;sup>2</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>() =</sup> Estimate generated from less than 5 unweighted observations

## Payment for family planning

Percentage of women ages 15 to 49 currently using a modern contraceptive method who paid fees for services, by marital status and background characteristics

<b>Background characteristics</b>	All women who used in past 12 months	All married women who used in past 12 month
Number	109	74
Total	51.5	52.0
Age group		
15–19	(53.7)	(74.2)
20–24	(53.2)	(65.1)
25–29	(64.3)	(69.1)
30–34	(53.9)	(55.6)
35–39	(48.9)	(43.2)
40–44	(39.6)	(39.6)
45–49	(13.2)	(13.2)
Marital status		
Married <sup>2</sup>	52.0	na
Unmarried, sexually active	(40.4)	na
Parity	(1211)	
0-1	48.3	(72.8)
2–3	(63.0)	(55.5)
4 or more	48.5	46.8
Residence		
Urban	57.8	(63.3)
Rural	48.4	47.7
Education		
Never attended	(59.1)	(52.4)
Primary	49.9	(48.1)
Secondary	49.1	(58.6)
Higher	(52.7)	(44.6)
Wealth quintile		
Lowest	(55.9)	(54.4)
Lower	(24.6)	(24.0)
Middle	(55.3)	(56.7)
Higher	(70.2)	(67.6)
Highest	(56.6)	(56.9)

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

<sup>&</sup>lt;sup>2</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>( ) =</sup> Estimate generated from less than 5 unweighted observations

#### **Method information index**

Percentage of current modern contraceptive users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics<sup>2</sup>

<b>Background characteristics</b>	othe	rmed about er methods	sid	med about e effects		hat to do if ed side effects
	All users	Married users	All users	Married users	All users	Married users
Number	109	74	85	56	32	25
Total	32.2	37.4	36.9	44.3	72.5	74.7
Age group						
15–19	(18.4)	(53.3)	(28.3)	(35.6)	(63.1)	(100.0)
20–24	(13.2)	(0.0)	(18.5)	(45.2)	(31.9)	(50.8)
25–29	(41.7)	(45.5)	(60.1)	(68.6)	(79.8)	(74.3)
30–34	(38.6)	(35.7)	(22.0)	(25.9)	(72.4)	(72.4)
35–39	(42.9)	(42.2)	(41.9)	(40.9)	(85.0)	(82.4)
40–44	(20.6)	(20.6)	(53.1)	(53.1)	(61.1)	(61.1)
45–49	(71.9)	(71.0)	(46.7)	(46.7)	(100.0)	(100.0)
Marital status	. ,	, ,	, ,			,
Married <sup>3</sup>	37.4	na	44.3	na	74.7	na
Unmarried, sexually active	(13.3)	na	(18.3)	na	(40.0)	na
Parity Parity	(13.3)	na	(10.5)	IIG	(70.0)	nu
0–1	16.8	(29.0)	21.7	(50.1)	(59.3)	(67.4)
2–3	(51.0)	(47.1)	(67.8)	(67.0)	(77.1)	(79.0)
4 or more	35.4	35.0	36.0	35.5	(75.3)	(73.9)
Residence	33.1	55.0	50.0	55.5	(13.5)	(15.5)
Urban	42.5	(62.2)	35.6	(55.1)	(90.2)	(90.2)
Rural	27.1	28.0	37.7	38.2	(62.7)	(61.8)
Education	27.11	20.0	57.7	20.2	(02.7)	(01.0)
Never attended	(34.3)	(35.3)	(25.3)	(24.7)	(66.4)	(81.0)
Primary	33.6	(33.5)	(47.8)	(47.7)	(80.1)	(78.4)
Secondary	26.2	(41.3)	33.3	(45.0)	(67.2)	(65.9)
Higher	(49.8)	(43.2)	(48.2)	(78.1)	(76.7)	(76.7)
	(12.0)	(13.2)	(10.2)	(70.1)	(70.7)	(70.7)
Wealth quintile	(14.0)	(16.7)	(25.0)	(20.6)	(100.0)	(100.0)
Lowest	(14.0)	(16.7)	(25.0)	(28.6)	(100.0)	(100.0)
Lower	(19.7)	(30.4)	(36.1)	(27.6)	(29.9)	(0.0)
Middle	(43.5)	(37.7)	(23.6)	(15.9)	(100.0)	(100.0)
Higher	(24.0)	(21.1)	(37.2)	(35.1)	(25.8)	(30.3)
Highest	43.1	(61.1)	44.2	(65.9)	(93.2)	(92.8)

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

- () = Estimate generated from less than 25 unweighted observations
- () = Estimate generated from less than 5 unweighted observations

<sup>&</sup>lt;sup>2</sup>Informed about other methods" asked of all current contraceptive users, "informed about side effects" asked of current modern contraceptive users, and "told what to do if experienced side effects" asked of current modern contraceptive users who indicated they were told about side effects.

<sup>&</sup>lt;sup>3</sup>Married includes married women and those who are living with a man

## Quality of care for family planning

Percentage of current modern contraceptive users who would return, would refer a friend/relative, or would return and refer friend/relative to provider, by background characteristics

Background characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Number	85	85	85
Total	79.1	78.2	75.9
Age group			
15–19	(75.4)	(75.4)	(75.4)
20–24	(87.0)	(81.7)	(81.7)
25–29	(75.0)	(75.0)	(75.0)
30–34	(78.6)	(71.4)	(67.0)
35–39	(84.3)	(84.3)	(84.3)
40–44	(66.9)	(79.4)	(66.9)
45–49	(100.0)	(100.0)	(100.0)
Marital status			
Married <sup>2</sup>	74.2	74.0	70.5
Unmarried,	(88.0)	(85.2)	(85.2)
sexually active			
Parity			
0–1	81.8	79.8	79.8
2–3	(84.2)	(89.1)	(84.2)
4 or more	74.7	72.3	69.2
Residence			
Urban	84.9	80.7	80.7
Rural	75.7	76.8	73.1
Education			
Never attended	(66.2)	(71.0)	(66.2)
Primary	(77.0)	(72.5)	(66.7)
Secondary	83.4	81.9	81.9
Higher	(88.8)	(88.8)	(88.8)
Wealth quintile			
Lowest	(44.8)	(58.0)	(44.8)
Lower	(89.8)	(84.5)	(84.5)
Middle	(75.7)	(75.7)	(75.7)
Higher	(72.3)	(77.9)	(72.3)
Highest	86.9	81.6	81.6

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

<sup>&</sup>lt;sup>2</sup>Married includes married women and those who are living with a man

na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>( ) =</sup> Estimate generated from less than 5 unweighted observations

# **Knowledge of family planning**

Percentage of women ages 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics

Background characteristics		alth worker who amily planning		lity and provider family planning	Either visited by health worker or visited facility and discussed family planning		
Number	All women <b>850</b>	Married women 592	All women <b>850</b>	Married women 592	All women <b>850</b>	Married women 592	
Total	9.6	10.9	11.9	15.3	17.6	21.5	
Age group							
15–19	5.5	6.3	4.4	7.9	7.9	10.8	
20–24	8.0	10.3	11.6	16.3	15.7	21.4	
25–29	7.9	8.0	17.2	19.3	21.0	23.9	
30–34	16.9	13.7	12.3	12.8	25.8	22.6	
35–39	14.3	16.9	22.3	23.5	28.8	31.6	
40–44	11.4	13.3	8.9	10.4	15.4	18.0	
45–49	3.8	4.7	9.8	12.0	11.9	14.5	
Marital status							
Married <sup>1</sup>	10.9	na	15.3	na	21.5	na	
Unmarried, sexually active	3.3	na	6.1	na	7.7	na	
Parity							
0-1	6.5	8.4	5.6	10.1	10.4	15.5	
2–3	12.1	10.4	15.8	16.2	22.8	22.7	
4 or more	11.7	12.7	17.4	17.9	23.3	24.3	
Residence							
Urban	10.6	12.2	10.5	19.6	19.6	29.0	
Rural	9.3	10.6	12.2	14.6	17.1	20.2	
Education							
Never attended	10.1	10.5	11.4	12.5	16.4	17.4	
Primary	8.7	10.0	15.4	18.3	21.0	25.7	
Secondary	8.8	12.2	10.7	18.2	15.8	24.3	
Higher	15.0	13.6	5.4	10.4	20.4	24.0	
Wealth quintile							
Lowest	8.2	8.6	11.6	13.8	16.7	19.3	
Lower	9.1	10.5	13.0	15.8	20.1	24.1	
Middle	8.6	10.3	11.5	12.6	15.5	17.8	
Higher	9.8	12.4	9.8	13.9	13.1	17.1	
Highest	11.5	12.4	13.3	20.4	21.7	28.9	

<sup>&</sup>lt;sup>1</sup>Married includes married women and those who are living with a man na = not applicable

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

# List of indicators for sampling errors, PMA2016/Taraba

Variable	Estimation	Base population
Currently using a modern method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using a traditional method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using any contraceptive method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using injectables	Proportion	All women ages 15–49 Married women ages 15–49
Currently using male condoms	Proportion	All women ages 15–49 Married women ages 15–49
Currently using implants	Proportion	All women ages 15–49 Married women ages 15–49
Chose method by self or jointly in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Paid fees for family planning services in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about other methods	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about side effects	Proportion	All women ages 15–49 Married women ages 15–49
Satisfied with provider: Would return and refer friend/relative to provider	Proportion	All women ages 15–49 Married women ages 15–49
Visited by health worker who talked about family planning information in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49

# **Characteristics of service delivery points**

# Number and percentage of service delivery points (SDPs) by background characteristics

<b>Background characteristics</b>	Number	Percentage
Total	49	100.0
SDP type		
Hospital	4	8.2
Health center <sup>1</sup>	21	42.9
Pharmacy <sup>2</sup>	21	42.9
Other	3	6.1
Sector		
Public	24	49.0
Private	25	51.0
Offers family planning		
No	8	16.3
Yes	41	83.7

<sup>&</sup>lt;sup>1</sup>Includes Health Post, Health Clinic, and Maternity Center

<sup>&</sup>lt;sup>2</sup>Includes Chemist and Patent Medicine Vendors

# **Charging fees for family planning**

Percentage of service delivery points (SDPs) that charge fees for family planning services, by sector and background characteristics, among SDPs offering family planning

Background characteristic	Sector						
	Public	Private					
Number of SDPs	20	21					
Total	(35.0)	(95.2)					
SDP type		, ,					
Hospital	(25.0)	na					
Health center <sup>1</sup>	(33.3)	(100.0)					
Pharmacy <sup>2</sup>	na	(94.7)					
Other	(100.0)	(100.0)					

<sup>&</sup>lt;sup>1</sup>Includes Health Post, Health Clinic, and Maternity Center

<sup>&</sup>lt;sup>2</sup>Includes Chemist and Patent Medicine Vendors

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>() =</sup> Estimate generated from less than 5 unweighted observations

## Contraceptive choice: Availability of at least 3 or 5 modern contraceptive methods

Percentage of public and private service delivery points (SDPs) offering at least 3 or 5 modern contraceptive methods<sup>1</sup>, by sector and background characteristics, among all SDPs

Background characteristic	3 or more	e methods	5 or more methods				
Number of SDPs	Public 24	Private 25	Public 24	Private 25			
Total	(54.2)	20.0	(54.2)	20.0			
SDP type							
Hospital	(100.0)	na	(100.0)	na			
Health Center <sup>1</sup>	(42.1)	(0.0)	(42.1)	(0.0)			
Pharmacy <sup>2</sup>	na	(23.8)	na	(23.8)			
Other	(100.0)	(0.0)	(100.0)	(0.0)			

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

<sup>&</sup>lt;sup>2</sup>Includes Health Post, Health Clinic, and Maternity Center

<sup>&</sup>lt;sup>3</sup>Includes Chemist and Patent Medicine Vendors

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>() =</sup> Estimate generated from less than 5 unweighted observations

## Contraceptive choice: Availability of modern contraception, by method

Percentage of service delivery points (SDPs) offering different types of contraceptive methods<sup>1</sup>, by sector and method, among all SDPs

	Public SDPs										
			Co	ontracepti	ve method	l					
SDP characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception				
Number = 24							•				
Total	(70.8)	(70.8)	(45.8)	(54.2)	(75.0)	(16.7)	(37.5)				
SDP type											
Hospital	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(75.0)	(100.0)				
Health center <sup>2</sup>	(63.2)	(63.2)	(31.6)	(42.1)	(68.4)	(0.0)	(21.1)				
Pharmacy <sup>3</sup>	na	na	na	na	na	na	na				
Other	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)				

	Private SDPs										
	Contraceptive method										
SDP characteristic	Pill Injectables		IUD	Implants	Male condom	Female sterilization	Emergency contraception				
Number = 25							•				
Total	72.0	36.0	4.0	8.0	64.0	0.0	8.0				
SDP type											
Hospital	na	na	na	na	na	na	na				
Health center <sup>2</sup>	(50.0)	(50.0)	(50.0)	(50.0)	(50.0)	(0.0)	(0.0)				
Pharmacy <sup>3</sup>	(76.2)	(33.3)	(0.0)	(0.0)	(66.7)	(0.0)	(9.5)				
Other	(50.0)	(50.0)	(0.0)	(50.0)	(50.0)	(0.0)	(0.0)				

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

<sup>&</sup>lt;sup>2</sup>Includes Health Post, Health Clinic, and Maternity Center

<sup>&</sup>lt;sup>3</sup>Includes Chemist and Patent Medicine Vendors

<sup>( ) =</sup> Estimate generated from less than 25 unweighted observations ( ) = Estimate generated from less than 5 unweighted observations

## **Contraceptive stock-outs, by method**

Percentage of service delivery points (SDPs) stocked out of specific modern contraceptives <sup>1</sup> at time of survey or stocked out in the last 3 months, by sector and background characteristics, among SDPs that offer each method

	Public SDPs									
	Contraceptive method									
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception				
Number of SDPs	17 17		11	13	18	9				
Total	(41.2)	(29.4)	(27.3)	(53.8)	(33.3)	(55.6)				
SDP type										
Hospital	(25.0)	(25.0)	(0.0)	(25.0)	(50.0)	(75.0)				
Health center <sup>2</sup>	(50.0)	(33.3)	(50.0)	(75.0)	30.769	(50.0)				
Pharmacy <sup>2</sup>	na	na	na	na	na	na				

	Private SDPs									
		(	Contrace	ptive meth	od					
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception				
Number of SDPs	18 9		1	2	16	2				
Total	(27.8)	(44.4)	(100.0)	(50.0)	(18.8)	(100.0)				
SDP type										
Hospital	na	na	na	na	na	na				
Health center <sup>2</sup>	(100.0)	(100.0)	(100.0)	(0.0)	(100.0)	na				
Pharmacy <sup>3</sup>	(18.8)	(28.6)	na	(0.0)	(14.3)	(100.0)				

<sup>&</sup>lt;sup>1</sup>Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

<sup>&</sup>lt;sup>2</sup>Includes Health Post, Health Clinic, and Maternity Center

<sup>&</sup>lt;sup>3</sup>Includes Chemist and Patent Medicine Vendors

<sup>() =</sup> Estimate generated from less than 25 unweighted observations

<sup>() =</sup> Estimate generated from less than 5 unweighted observations

# Number of family planning visits, by method

Total number of visits and new clients for contraceptive services, by method type and background characteristics, among advanced service delivery points (SDPs) that offer each method

	Contraceptive Methods													
	Steriliz	zation	Injectable		IUD		Implants		Male condom		Pills		Emergency Contraception	
	Total women	Total men	Total	New	Total	New	Total	New	Total	New	Total	New	Total	New
Number of SDPs	4	4	19	)	12	2		15	2	0	19	)	ç	9
Total	65	3	1,466	359	226	53	921	224	1,470	652	1,423	402	530	78
SDP characteristic														
SDP type														
Hospital	30	1	533	109	110	22	336	94	999	471	618	304	80	32
Health center <sup>2</sup>	na	na	223	46	16	1	80	29	385	156	304	95	50	6
Other	35	2	710	204	100	30	505	101	86	25	501	3	400	40
Sector														
Public	65	3	1,451	355	223	53	914	223	1,464	652	1,422	401	530	78
Private	na	na	15	4	3	0	7	1	6	0	1	1	0	0

<sup>&</sup>lt;sup>1</sup>Hospital, Health Clinic, and other facilities

<sup>&</sup>lt;sup>2</sup>Includes Health Post, Health Clinic, and Maternity Center