

PMA ABORTION SURVEY RESULTS : BURKINA FASO

December 2020 - March 2021



KEY FINDINGS:

Abortion is a common reproductive health event in Burkina Faso, but more than half are unsafe, involving non-recommended methods from non-clinical sources.



Nearly three in ten women who reported an abortion indicated experiencing a potential severe complication, and only half of these women accessed postabortion care at a facility for treatment.



75% of women were unaware that abortion could be accessed safely in certain situations under the law in Burkina Faso.

ABORTION IN BURKINA FASO: LEGALLY RESTRICTED BUT NOT UNCOMMON

Induced abortion is legally permitted in Burkina Faso in cases of rape, incest, fetal defects, or when the woman's life or physical health are in danger. Approximately 25 abortions per 1,000 women were done in Burkina Faso in 2008, of which 43% resulted in complications, according to estimates extrapolated from postabortion care (PAC) data.¹ While these estimates provide an overall assessment of the extent of abortion in Burkina Faso and the level of risk associated, we have limited information about who experiences abortion, and, in particular, those who use procedures associated with a heightened risk of morbidity and mortality.

THE PMA BURKINA FASO ABORTION STUDY

Between December 2020 and March 2021, Performance Monitoring for Action (PMA) conducted a population-based survey to produce updated national estimates of abortion indicators in Burkina Faso and to understand the associated health risks. Details regarding the methodology of this study are described at the end of this brief.

ABORTION IS COMMON, BUT VARIES ACROSS A WOMAN'S LIFESPAN

Overall, the annual abortion incidence in Burkina Faso in 2020² is estimated at 23 abortions per 1,000 women aged 15 to 49, equivalent to approximately 113,000 abortions. The abortion incidence rate is higher among younger, unmarried and nulliparous women. Likewise, the incidence rate was higher among more educated, and urban women.


ANNUAL ABORTION INCIDENCE IN BURKINA FASO

Annual incidence of abortion in Burkina Faso, by sociodemographic characteristics of women, 2020 (n=6,388)



PRIMARY REASONS FOR ABORTION IN BURKINA FASO

Reasons for abortion varied across the lifespan, but are often related to early pregnancies that upset social norms and interfere with education.




“Because for a while I thought the world was crumbling on me. I had thought that everything I did for my life was destroyed. And with the child, I wasn’t going to be able to be who I am anymore and that I had failed everything. So I had an abortion.”

Single woman, age 32 with 4 children at time of abortion

THE MAJORITY OF ABORTIONS ARE UNSAFE, WHICH INCREASES COMPLICATIONS

Over half of abortions were most unsafe (51%)³, involving a non-clinical provider (i.e. not a public or private health facility) and a non-recommended method (i.e. something other than abortion surgery or misoprostol with or without mifepristone). Unsafe abortions are most common among women with no education, unmarried women, and the oldest women.

DISTRIBUTION OF LEVEL OF ABORTION SAFETY ⁴



PMA DEFINITIONS OF ABORTION SAFETY

Abortion safety was operationalized into four categories using abortion method and source data as follows:

1. Recommended method(s) (i.e. surgery or mifepristone/misoprostol) from clinical source(s) (i.e. public or private healthcare facilities)
2. Recommended method(s) involving non-clinical source(s)
3. Non-recommended method(s) from clinical source(s)
4. Non-recommended method(s) involving non-clinical source(s)

Abortions in the fourth category were deemed most unsafe.


**“Interviewer: Did the old lady explain it to you? Did she tell you how it goes?
Respondent: That she, she gives you the herbal tea to drink and that during the three days, the belly will hurt really bad; that the belly will hurt very bad and that when it hurts that to take products to calm; that’s what didn’t motivate me.”**

Single woman, age 15 and no children at time of abortion


“The care was not good because with all these products I was gorging myself with, I was even in danger of losing my life. With all these... how shall I put it? These plants that I was introducing (into my vagina) I could even maybe give myself infections or even cause other diseases.”

Single woman, age 19 with no children at time of abortion

ABORTION METHOD




ABORTION FACILITY




PERCENT OF ABORTIONS IN BURKINA FASO CONSIDERED UNSAFE BY RESPONDENTS' BACKGROUND CHARACTERISTICS


AGE




EDUCATION




CURRENTLY MARRIED



RESIDENCE



PARITY



COMPLICATIONS AND ACCESS TO POSTABORTION CARE



of women reported a potential severe complication (fever, vaginal discharge, punctured uterus or other complication requiring surgery); this proportion increases to 32% among those reporting most unsafe abortion.

Only half of women who described potentially severe complications accessed facilities for postabortion care.

“Very! Very heavy. In the meantime, I didn’t even have the strength anymore and every drop that fell, before the drop itself fell, you cry. You cry before the drop itself falls and it’s as if pieces of your very flesh are coming out. Every time you bleed, you cry, you cry, it wasn’t easy.”

Single woman, age 22 with no child at time of abortion

“If you have an abortion, even if it’s done badly, after if you go to the hospital to get treatment it won’t be easy. So that you have the care. You’re going to have in any case, but maybe there will be complications even.”

Single woman, age 22 with 1 child at time of abortion


WOMEN NEED ACCURATE INFORMATION ABOUT ABORTION

Many women were unaware of the legal ground for abortion in Burkina Faso and the majority were unaware of safe abortion methods.

Knowledge

“We are not in Europe. There is no law that says we accept that the woman has an abortion if she doesn’t want to. I learned that there is that in one country but not in Burkina, so it’s difficult, so the care is complicated.”

Single woman, age 21 with no children at time of abortion



of women were unaware that abortion could be accessed safely under the law in cases of rape, incest, fetal impairment, or when the woman’s life or physical health are in danger.


Only 7% of women knew any recommended abortion method (i.e., surgery or medication abortion pills), most of whom only knew about surgery (6%).

Women with no education, women with the least financial resources, and women residing in rural areas were less knowledgeable about the law and safe abortion methods.

Decision-making

39% of women in Burkina Faso would recommend a friend go to a public health facility if she needed to terminate a pregnancy.

REASONS FOR CHOOSING ABORTION METHOD




¹ Sedgh G, Rossier C, Kaboré I, Bankole A, Mikulich M. Estimating abortion incidence in Burkina Faso using two methodologies. *Studies in Family Planning*. 2011;42(3):147-154.

² Abortion incidence estimates come from adjusted friend data.

³ Abortion safety estimates from respondent data.

⁴ Estimates are similar to those for confidantes.

ACCESSING ABORTION CARE IS CHALLENGING FOR MANY WOMEN



of women said it was very or somewhat difficult to pay for their abortion. Women with the least financial resources were more than twice as likely to report having difficulty paying for an abortion (55%).



said they paid a bribe to the abortion provider, and this proportion increases to 59% among those who accessed care at a private health facility.

“In all sincerity the price was too much, I can say, because it was not easy for us; we had to look left and right, take out loans with people to be able to do it and then we paid them back.”

Married woman, age 36 with 2 children at time of abortion

RECOMMENDATIONS

Findings indicate that women often rely on abortion – most often under unsafe conditions – to manage their fertility in the context of experiencing an unwanted pregnancy. The data show social inequities in access to information and care, with more disadvantaged women having less information about recommended abortion methods and more likely to rely on unsafe methods and non-clinical providers. The Burkina Faso Ministry of Health, non-governmental organizations (NGOs), and civil society working in the reproductive health field can take the following actions to reduce the burden of unsafe abortion and associated negative impacts on maternal health:

- Increase information about the availability of quality, voluntary family planning services and improve access to contraceptive methods throughout the health system to prevent unintended pregnancies.
- Ensure the availability of safe abortion and postabortion care services to the full extent of the law, particularly at primary care facilities that serve the most disadvantaged populations. The availability of care requires the continuation and expansion of training for doctors, midwives and nurses, as well as ensuring the supply of materials and medications such as manual vacuum aspiration kits and mifepristone and misoprostol pills.
- Inform the public about the specific conditions under which abortion is considered legal in Burkina Faso.
- Inform communities and women about contraception, risks associated with abortion using non-recommended methods, and safe abortion and postabortion care services and the locations where these services can be availed.

Taken together, these changes can significantly reduce the extent of unsafe abortion, associated complications, and disparities, and reduce the hundreds of preventable unsafe abortion-related maternal deaths that occur each year in Burkina Faso.

What is PMA?

PMA uses mobile technology and female resident data collectors to support rapid-turnaround surveys to monitor key family planning and health indicators in Africa and Asia. PMA Burkina Faso is led by l'Institut Supérieur des Sciences de la Population at l'Université Joseph Ki-Zerbo, Ougadougou, Burkina Faso. Overall direction and support are provided by the Bill & Melinda Gates Institute for Population and Reproductive Health at the Johns Hopkins University and Jhpiego. Funding for PMA is provided by the Bill & Melinda Gates Foundation; an Anonymous Donor provided funding for the abortion module.

PMA Burkina Faso collected information on knowledge, practice, and coverage of family planning services in 167 enumeration areas selected using a multi-stage stratified cluster design with urban-rural strata. The results are representative at the national level and the urban/rural areas. Phase 2 data were collected from December 2020 and March 2021 from 5,522 households (97.9% response rate) and 6,388 women 15-49 years old (93.4% response rate). For further sampling information and full datasets, visit <https://www.pmadata.org/countries/burkina-faso>. For this phase of data collection, we added an abortion module to estimate abortion incidence and safety. This included asking questions about the respondent's experience with abortion, as well as her closest friend's. Details on the friend abortion incidence measurement methodology and our safety measurement approach are provided elsewhere [Bell, S. O., M. Shankar, E. Omoluabi, A. Khanna, H. K. Andoh, F. OlaOlorun, D. Ahmad, G. Guiella, S. Ahmed and C. Moreau (2020). "Social network-based measurement of abortion incidence: promising findings from population-based surveys in Nigeria, Cote d'Ivoire, and Rajasthan, India." *Population Health Metrics* 18(1): 1-15; Bell, S. O., F. OlaOlorun, M. Shankar, D. Ahmad, G. Guiella, E. Omoluabi, A. Khanna, A. K. Hyacinthe and C. Moreau (2019). "Measurement of Abortion Safety Using Community-Based Surveys: Findings from Three Countries." *PLoS One* 14(11): 1-14.; Bell, S. O., E. Omoluabi, F. OlaOlorun, M. Shankar and C. Moreau (2020). "Inequities in the incidence and safety of abortion in Nigeria." *BMJ Global Health* 5(1): e001814.]. Data collectors also followed up with and conducted in-depth qualitative interviews with 30 women who reported an abortion in the PMA study and consented to be recontacted.